arXiv:math/0509154v1 [math.AG] 7 Sep 2005

Irreducibility of Hurwitz spaces

Vassil Kanev

Abstract

Graber, Harris and Starr proved, when $n \geq 2d$, the irreducibility of the Hurwitz space $\mathcal{H}^0_{d,n}(Y)$ which parametrizes degree d coverings of a smooth, projective curve Y of genus ≥ 1, simply branched in n points, with full monodromy group S_d. We sharpen this result and prove that $\mathcal{H}^0_{d,n}(Y)$ is irreducible if $n \geq \max\{2, 2d-4\}$ and in the case of elliptic Y if $n \geq \max\{2, 2d-6\}$. We extend the result to coverings simply branched in all but one point of the discriminant. Fixing the ramification multiplicities over the special point we prove that the corresponding Hurwitz space is irreducible if the number of simply branched points is $\geq 2d-2$. We study also simply branched coverings with monodromy group $\neq S_d$ and when n is large enough determine the corresponding connected components of $\mathcal{H}_{d,n}(Y)$. Our results are based on explicit calculation of the braid moves associated with the standard generators of the n-strand braid group of Y.

Introduction

Let Y be a smooth, connected, projective complex curve of genus $g \geq 0$. Let $\mathcal{H}_{d,n}(Y)$ be the Hurwitz space which parametrizes degree d coverings of Y simply branched in n points. A classical result of Hurwitz [Hu] states that $\mathcal{H}_{d,n}(\mathbb{P}^1)$ is irreducible. More generally one can consider coverings of $\mathbb{P}^1$ which are simply branched in all but one point of the discriminant. Fixing the ramification multiplicities over the special point one obtains a corresponding Hurwitz space which turns out to be irreducible as well (see [Na], [Kl], [Mo]).

Coverings of curves of positive genus were studied by Graber, Harris and Starr in [GHS]. They considered the Hurwitz space $\mathcal{H}^0_{d,n}(Y)$ which parametrizes coverings with full monodromy group S_d and proved it is irreducible if $n \geq 2d$. Another result of this type was obtained by F. Vetro [Ve].

In the present paper we sharpen the result of Graber, Harris and Starr and prove that $\mathcal{H}^0_{d,n}(Y)$ is irreducible if $n \geq 2d-2$ (cf. Theorem 3.10). Our approach allows to extend the result to coverings which are simply branched in all but one point of the discriminant. Fixing the branching data of the special point, i.e. a partition $\underline{e} = \{e_1, e_2, \ldots, e_q\}$ where

---

Preprint N. 241, Dipartimento di Matematica, Università di Palermo.

2000 *Mathematics Subject Classification.* Primary: 14H10; Secondary: 14H30, 20F36.

$e_1 \geq e_2 \geq \ldots \geq e_q \geq 1$ and $e_1 + \cdots + e_q = d$ one obtains a Hurwitz space $\mathcal{H}^0_{d,n,\underline{e}}(Y)$ parametrizing coverings with full monodromy group, simply branched in n points and ramified with multiplicities $e_1, \ldots, e_q$ over one additional point. We prove $\mathcal{H}^0_{d,n,\underline{e}}(Y)$ is irreducible if $n \geq 2d - 2$ (cf. Theorem 3.11). Coverings with $n < 2d - 2$ are more difficult to deal with. We prove the irreducibility of $\mathcal{H}^0_{d,n}(Y)$ when $n = 2d - 4,\ d \geq 3,\ g(Y) \geq 1$ (cf. Theorem 4.5) and the irreducibility of $\mathcal{H}^0_{d,n}(Y)$ when $n = 2d - 6,\ d \geq 4,\ g(Y) = 1$ (cf. Theorem 5.4). The case of simply branched coverings with monodromy groups $\neq S_d$ can be reduced via étale coverings to the case of coverings with full monodromy group. In Theorem 3.14 and Theorem 5.5 we prove that connected components of $\mathcal{H}_{d,n}(Y)$ not contained in $\mathcal{H}^0_{d,n}(Y)$ may exist only if d is not prime and in this case if n is sufficiently large such connected components correspond bijectively to the equivalence classes of étale coverings $[\tilde{Y} \to Y]$ of degree d_2, where $d_2|d,\ d_2 \neq 1, d$.

In proving these results we follow the standard approach for determining the connected components of $\mathcal{H}_{d,n}(Y)$. Associating to every equivalence class of coverings $[X \to Y] \in \mathcal{H}_{d,n}(Y)$ its discriminant locus yields an étale mapping $\mathcal{H}_{d,n}(Y) \to Y^{(n)} - \Delta$. Fixing a $D \in Y^{(n)} - \Delta$ the fiber over D is identified via Riemann's existence theorem to the equivalence classes modulo inner automorphisms of ordered $(n+2g)$-tuples $(t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$ satisfying $t_1 \cdots t_n = [\lambda_1, \mu_1] \cdots [\lambda_g, \mu_g]$ – we call such $(n+2g)$-tuples *Hurwitz systems* – where $t_i, \lambda_k, \mu_k \in S_d$, t_i are transpositions, and t_i, λ_k, μ_k with $1 \leq i \leq n,\ 1 \leq k \leq g$ generate a transitive subgroup of S_d. The problem is thus reduced to finding the orbits of the action of the n-strand braid group of Y, namely $\pi_1(Y^{(n)} - \Delta, D)$, on the set of equivalence classes of Hurwitz systems. In fact it is more natural and it suffices to study the action of the braid group of the open Riemann surface $Y - \{b_0\}$, where b_0 is a fixed point. Birman found in [Bi] a natural system of generators of these braid groups. Our results are based on Theorem 1.8 where we calculate explicitly how these generators act on Hurwitz systems.

We follow the key idea of [GHS] which consists in applying a sequence of braid moves to a given Hurwitz system $(t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$ in order to replace it by a new one $(\tilde{t}_1, \ldots, \tilde{t}_n; \tilde{\lambda}_1, \tilde{\mu}_1, \ldots, \tilde{\lambda}_g, \tilde{\mu}_g)$ in which as many as possible of the elements $\tilde{\lambda}_k, \tilde{\mu}_k$ equal 1 and then reduce the sequence $(\tilde{t}_1 \ldots, \tilde{t}_n)$ to a normal form using only elementary transformations of the Artin's braid group. An important tool in our arguments is Main Lemma 2.1 which states that if in a Hurwitz system $(t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$ one has $t_i t_{i+1} = 1$, then replacing the pair (t_i, t_{i+1}) by a pair $(h^{-1} t_i h, h^{-1} t_{i+1} h)$, where h belongs to the subgroup generated by t_j, λ_k, μ_k with $1 \leq j \leq n,\ j \neq i, i+1,\ 1 \leq k \leq g$, one obtains a braid-equivalent Hurwitz system, i.e. can be obtained from the initial one by a finite sequence of braid moves. This is a generalization of Lemma 2.2 of [GHS] proved there using the Kontsevich moduli space of stable maps. The proof of our Main Lemma 2.1 is more elementary and uses only the explicit formulas for the braid moves of Theorem 1.8. We think Theorem 1.8 and Main Lemma 2.1 are results of independent interest, since both are valid for Hurwitz systems with values in an arbitrary (possibly infinite) group G. We profitted a lot from the paper of Mochizuki [Mo] which was the starting point for us in studying the problem of the irreducibility of Hurwitz spaces.

Notation and conventions. We assume that the base field $k = \mathbb{C}$. Throughout the paper

$\pi : X \to Y$ denotes a covering, i.e. a finite holomorphic mapping, of smooth projective curves. We do not assume that X is connected, while Y is always assumed connected. Two coverings $\pi_1 : X_1 \to Y$ and $\pi_2 : X_2 \to Y$ are called equivalent if there is an isomorphism $f : X_1 \to X_2$ such that $\pi_1 = \pi_2 \circ f$. A covering $\pi : X \to Y$ of degree d is called simple if both X and Y are irreducible and for each $y \in Y$ one has $d - 1 \leq \# \, \pi^{-1}(y) \leq d$. We use mainly right actions and write x^g for $x \in \Sigma$ and $g \in G$, where the group G acts on the set Σ on the right. In particular if $\Sigma = G$ we let $x^g = g^{-1}xg$.

Contents. 1. Braid moves, 2. The main lemma, 3. The case $n \geq 2d-2$, 4. The case $n = 2d-4$, $g \geq 1$, 5. The case $n = 2d-6$, $g = 1$.

1 Braid moves

1.1. The equivalence classes of (possibly nonconnected) coverings of degree d of a smooth, projective, irreducible curve Y branched in $n > 0$ points are parametrized by a smooth scheme $\mathcal{H}_d^{(n)}(Y)$ which is an étale cover of $Y^{(n)} - \Delta$. Here Δ is the codimension one subscheme consisting of nonsimple divisors of degree n. The covering mapping associates to $[X \to Y]$ its branch locus D. Let $b_0 \in Y$. We denote by $\mathcal{U}(b_0)$ the open subset of $\mathcal{H}_d^{(n)}(Y)$ consisting of coverings which are unramified at b_0. Similarly we consider equivalence classes of pairs $[\pi : X \to Y, \phi]$ where $[X \to Y] \in \mathcal{U}(b_0)$ and $\phi : \pi^{-1}(b_0) \to \{1, \ldots, d\}$ is a bijection. We denote by $\mathcal{H}_d^{(n)}(Y, b_0)$ the corresponding étale cover of $\mathcal{U}(b_0)$. Let $D \in Y^{(n)} - \Delta$, $b_0 \notin D$. Riemann's existence theorem (see e.g. [Fu] p.544) establishes the following natural one-to-one correspondences

A. For $\mathcal{H}_d^{(n)}(Y, b_0)$ between:

  (i) the fiber of $\mathcal{H}_d^{(n)}(Y, b_0) \to (Y - b_0)^{(n)} - \Delta$ over D

  (ii) the set of homomorphisms $m : \pi_1(Y - D, b_0) \to S_d$ which satisfy the condition that $m(\gamma) \neq 1$ for each small loop which circles a point of D.

B. For $\mathcal{H}_d^{(n)}(Y)$ between:

  (i) the fiber of $\mathcal{H}_d^{(n)}(Y) \to Y^{(n)} - \Delta$ over D

  (ii) the set of equivalence classes modulo inner automorphisms of homomorphisms $m : \pi_1(Y - D, b_0) \to S_d$ satisfying the condition of A(ii), here $m \sim m'$ if there exists an $s \in S_d$ such that $m' = s^{-1}ms$.

We fix our preferences in this paper as follows. We consider the product in a fundamental group induced by product of arcs defined as

$$\gamma_1 * \gamma_2 = \begin{cases} \gamma_1(2t) & \text{for } 0 \leq t \leq 1/2, \\ \gamma_2(2t-1) & \text{for } 1/2 \leq t \leq 1 \end{cases} \tag{1}$$

We consider right action of S_d on $\{1,\dots,d\}$, so a permutation $\sigma \in S_d$ transforms i into i^σ and for instance $(123) = (12)(13)$. Explicitly the correspondence in (A) is defined in the following way. Given a pair $\pi : X \to Y, \quad \phi : \pi^{-1}(b_0) \to \{1,\dots,d\}$ let us number the elements of the fiber $\pi^{-1}(b_0)$ using ϕ, thus $\pi^{-1}(b_0) = \{x_1,\dots,x_d\}$ where $x_i = \phi^{-1}(i)$. Then given a homotopy class of loops $[\gamma]$ with $\gamma(0) = \gamma(1) = b_0$ the permutation $\sigma = m(\gamma)$ acts as follows. If lifting γ one starts from x_i and ends at x_j, then one lets $i^\sigma = j$. Let $s \in S_d$. Renumbering the points of the fiber $\pi^{-1}(b_0)$ by a bijective map $\psi : \pi^{-1}(b_0) \to \{1,\dots,d\}$ such that $\psi(x) = \phi(x)^s$ results in replacing the monodromy homomorphism m by $s^{-1}ms$. Hence the one-to-one correspondence in (B) is obtained from that of (A) applying the forgetful mapping $[X \to Y,\ \phi] \mapsto [X \to Y]$.

1.2. Suppose that $g(Y) \geq 1$. Let us fix the orientation of the real 2-manifold Y considered as a complex manifold. Let $D = \{b_1,\dots,b_n\} \subset Y$. Let $b_0 \notin D$. We describe a standard way of choosing generators of $\pi_1(Y - D, b_0)$. Cutting Y along $2g$ simple closed arcs which begin at b_0 and do not contain any of $b_i,\ i \geq 1$ one obtains a standard $4g$-polygon with sides $\alpha_1, \beta_1, \alpha_1^{-1}, \beta_1^{-1}, \dots, \alpha_g, \beta_g, \alpha_g^{-1}, \beta_g^{-1}$ which circle the polygon in counterclockwise direction. We consider a simple closed arc L which begins at b_0, $\quad L - b_0$ is contained in the interior of the 4g-polygon and passes consecutively in counterclockwise direction through the points $\{b_1,\dots,b_n\}$. The closed arc L divides the $4g$ polygon into two regions R and R' which stay on the left respectively on the right side of L with respect to its counterclockwise orientation. We choose a simple arc ℓ_1 which lies inside the region R and connects b_0 and b_1. Then we choose a second simple arc ℓ_2 inside R which connects b_0 with b_2, has only b_0 as point in common with ℓ_1, and lies on the left side of ℓ_1. Continuing in this way we obtain an ordered n-tuple $(\ell_1 \dots, \ell_n)$ of simple arcs which do not meet outside b_0. Let γ_i be a closed path which begins at b_0, travels along ℓ_i to a point near b_i, makes a small counterclockwise loop around b_i, and returns to b_0 along ℓ_i. We obtain a $(n+2g)$-tuple of closed arcs $(\gamma_1,\dots,\gamma_n; \alpha_1, \beta_1, \dots, \alpha_g, \beta_g)$ which we call a *standard system of closed arcs.* The corresponding homotopy classes yield a standard system of generators for $\pi_1(Y - D, b_0)$ which satisfy the only relation

$$\gamma_1\gamma_2\cdots\gamma_n \ \simeq \ [\alpha_1,\beta_1]\cdots[\alpha_g,\beta_g] \tag{2}$$

Figure 1 illustrates such a standard system. A reader who prefers the clockwise orientation of closed arcs and ordering of the branch points from left to right may look at this and all subsequent figures from the other side of the sheet. Given a covering $\pi : X \to Y$ with discriminant D and an isomorphism $\phi : \pi^{-1}(b_0) \to S_d$ one applies the monodromy homomorphism m and obtains $t_i = m(\gamma_i),\ \lambda_k = m(\alpha_k),\ \mu_k = \lambda_{g+k} = m(\beta_k)$.

Definition 1. An ordered sequence $(t_1,\dots,t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ of permutations in S_d with $t_i \neq 1$ for $\forall i$ and satisfying the relation

$$t_1t_2\cdots t_n \ = \ [\lambda_1,\mu_1]\cdots[\lambda_g,\mu_g]$$

is called a *Hurwitz system.* We let $\lambda_{g+k} = \mu_k$. We call the subgroup $G \subseteq S_d$ generated by all t_i, λ_k, μ_k the *monodromy group* of the Hurwitz system.



Figure 1: Standard system of closed arcs

By 1.1 given $D \in Y^{(n)} - \Delta$ and $b_0 \notin D$, and fixing the closed arcs $(\gamma_1, \ldots, \gamma_n; \alpha_1, \beta_1, \ldots, \alpha_g, \beta_g)$ as above, the fiber of $\mathcal{H}_d^{(n)}(Y, b_0) \to (Y - b_0)^{(n)} - \Delta$ over D may be identified with the set of all Hurwitz systems, while the fiber of $\mathcal{H}_d^{(n)}(Y) \to Y^{(n)} - \Delta$ over D may be identified with the set of equivalence classes $[t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g]$ of Hurwitz systems modulo inner automorphisms of S_d, where $(t_1, \ldots, \mu_g)$ is equivalent to $(t'_1, \ldots, \mu'_g)$ if there exists an $s \in S_d$ such that $t'_i = s^{-1} t_i s,\ \lambda'_k = s^{-1} \lambda_k s,\ \mu'_k = s^{-1} \mu_k s$ for $\forall i, k$.

An equivalent way of constructing a standard system of closed arcs is the following. One chooses first the $2g$ simple closed arcs $\alpha_1, \beta_1, \ldots, \alpha_g, \beta_g$. Then one chooses n simple arcs which start at b_0, lie inside the $4g$-polygon, do not meet outside b_0, and have for end points the n points of D. One enumerates these arcs according to the directions of departure in counterclockwise order. The obtained $(n + 2g)$-tuple $(\ell_1, \ldots, \ell_n; \alpha_1, \beta_1, \ldots, \alpha_g, \beta_g)$ is called an *arc system* (cf. [Lo] p.416). One considers the induced ordering of the points of D. One can take for R a star-like region which contains the union $\ell_1 \cup \ldots \cup \ell_n$ and let $L = \partial R$. In this way one obtains all ingredients used to construct a standard system of closed arcs $(\gamma_1, \ldots, \gamma_n; \alpha_1, \beta_1, \ldots, \alpha_g, \beta_g)$.

1.3. The connected components of $\mathcal{H}_d^{(n)}(Y)$ are in one-to-one correspondence with the orbits of the full n-strand braid group $\pi_1(Y^{(n)} - \Delta, D)$ acting on the fiber of the topological covering $\mathcal{H}_d^{(n)}(Y) \to Y^{(n)} - \Delta$ over D. Similar statement holds about $\mathcal{H}_d^{(n)}(Y, b_0)$ where one considers the braid group $\pi_1((Y - b_0)^{(n)} - \Delta, D)$. The identification of these fibers with the Hurwitz systems reduces the problem of determining the connected components of $\mathcal{H}_d^{(n)}(Y, b_0)$ and $\mathcal{H}_d^{(n)}(Y)$ to calculating the action of the respective braid groups on Hurwitz systems and then finding the orbits.

Let $\gamma_i, \alpha_k, \beta_k,\ 1 \le i \le n,\ 1 \le k \le g$ be a standard system of closed arcs obtained from an arc system $\ell_i, \alpha_k, \beta_k,\ 1 \le i \le n,\ 1 \le k \le g$ as in 1.2. Let $D^u,\ 0 \le u \le 1$ be a closed arc in $(Y - b_0)^{(n)} - \Delta$ with $D^0 = D^1 = D$. Suppose that starting from the given arc system one can extend the map $u \mapsto D^u$ to a homotopy of arc systems $\ell_i^u, \alpha_k^u, \beta_k^u,\ 1 \le i \le n, 1 \le k \le g$ based at b_0. This yields a corresponding homotopy of closed arcs $\gamma_i^u, \alpha_k^u, \beta_k^u\ 1 \le i \le n,\ 1 \le k \le g$ which form a standard system for each $u \in [0,1]$. Let $[X \to Y,\ \phi] \in \mathcal{H}_d^{(n)}(Y, b_0)$ have discriminant D and let $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ be the corresponding Hurwitz system. Then the lifting of the closed arc D^u, $0 \le u \le 1$, starting from a $[X \to Y,\ \phi]$, is defined by by $m^u : \pi_1(Y - D^u, b_0) \to S_d$ where (cf. [Fu] p.545)

$$m^u(\gamma_i^u) = t_i, \quad m^u(\alpha_k^u) = \lambda_k, \quad m^u(\beta_k^u) = \mu_k$$

Letting $\gamma_i^1 = \gamma_i',\ \alpha_k^1 = \alpha_k',\ \beta_k^1 = \beta_k'$ and $m^1 = m'$ we obtain the end point of the lifting of D^u is $[X' \to Y,\ \phi']$ whose monodromy map $m' : \pi_1(Y - D, b_0) \to S_d$ is defined by

$$m'(\gamma_i') = t_i, \quad m'(\alpha_k') = \lambda_k, \quad m'(\beta_k') = \mu_k$$

Evaluating m' at $\gamma_i, \alpha_k, \beta_k$ we obtain

$$m'(\gamma_i) = t_i', \quad m'(\alpha_k) = \lambda_k', \quad m'(\beta_k) = \mu_k'. \tag{3}$$

So $(t_1', \dots, t_n'; \lambda_1', \mu_1', \dots, \lambda_g', \mu_g')$ is the Hurwitz system corresponding to $[X' \to Y,\ \phi']$. Another approach is to consider

$$m(\gamma_i') = t_i'', \quad m(\alpha_k') = \lambda_k'', \quad m(\beta_k') = \mu_k''. \tag{4}$$

The Hurwitz system $(t_1'', \dots, t_n''; \lambda_1'', \mu_1'', \dots, \lambda_g'', \mu_g'')$ corresponds to a pair $[X'' \to Y,\ \phi'']$ whose monodromy map $m'' : \pi_1(Y - D, b_0) \to S_d$ satisfies

$$m''(\gamma_i) = t_i'', \quad m''(\alpha_k) = \lambda_k'', \quad m''(\beta_k) = \mu_k'' \tag{5}$$

Lemma 1.4. *Let us lift the path D^{1-u}, $0 \le u \le 1$ starting from $[X \to Y,\ \phi]$. Then the end point is $[X'' \to Y,\ \phi'']$.*

Proof. In terms of $\gamma_i', \alpha_k', \beta_k'$ the monodromy map of $[X \to Y,\ \phi]$ is given by $m(\gamma_i') = t_i'',\ m(\alpha_k') = \lambda_k'',\ m(\beta_k') = \mu_k''$. Let us consider the homotopy of closed paths $\gamma_i^{1-u}, \alpha_k^{1-u}, \beta_k^{1-u}$, $0 \le u \le 1$ and the path in $\mathcal{H}_d^{(n)}(Y, b_0)$ given by $n^u : \pi_1(Y - D^{1-u}, b_0) \to S_d$ where $n^u(\gamma_i^{1-u}) = t_i''$, $n^u(\alpha_k^{1-u}) = \lambda_k''$, $n^u(\beta_k^{1-u}) = \mu_k''$. Then $n^0 = m,\ n^1 = m''$ (cf. Eq.(5)) ☐

Definition 2. Given a closed arc D^u, $0 \le u \le 1$ in the configuration space $(Y - b_0)^{(n)} - \Delta$ the transformation of Hurwitz systems $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g) \mapsto (t_1', \dots, t_n'; \lambda_1', \mu_1', \dots, \lambda_g', \mu_g')$ given by Eq.(3) is called a *braid move* of the first type. The transformation $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g) \mapsto (t_1'', \dots, t_n''; \lambda_1'', \mu_1'', \dots, \lambda_g'', \mu_g'')$ given by Eq.(5) is called a braid move of the second type.

The braid moves of the first and of the second type are inverse to each other according to Lemma 1.4. It is evident that the braid moves of both types commute with inner automorphisms of S_d. So the braid moves are well-defined on equivalence classes of Hurwitz systems (cf. 1.2).

1.5. There is a convenient system of generators of $\pi_1((Y-b_0)^{(n)}-\Delta, D)$. We include here some material borrowed from [Bi] and [Sc] for the sake of convenience of the reader and since our choices differ slightly from theirs. Consider the Galois covering $p:(Y-b_0)^n \to (Y-b_0)^{(n)}$ with Galois group S_n. Restricting to the complement of Δ one obtains an unramified Galois covering $p:(Y-b_0)^n-p^{-1}(\Delta) \to (Y-b_0)^{(n)}-\Delta$. Following the notation of [FN] if $Q_1=\{b_0\}$ one denotes $(Y-b_0)^n-p^{-1}(\Delta)$ by $F_{1,n}Y$. Let $D=\{b_1,\dots,b_n\}$, $\tilde{D}=(b_1,\dots,b_n)$. One has an exact sequence

$$1 \longrightarrow \pi_1(F_{1,n}Y,\tilde{D}) \longrightarrow \pi_1((Y-b_0)^{(n)}-\Delta, D) \longrightarrow S_n \longrightarrow 1 \tag{6}$$

One determines first a system of generators of the pure braid group $\pi_1(F_{1,n}Y,\tilde{D})$ as follows. Consider the closed arcs in $F_{1,n}Y$ defined by $(b_1,\dots,b_{i-1},r_{ik}(t),b_{i+1},\dots,b_n)$ and $(b_1,\dots,b_{i-1},t_{ik}(t),b_{i+1},\dots,b_n)$, with $t\in[0,1]$, where r_{ik} and t_{ik} are the closed simple arcs based at b_i and pictured on Figure 2 by a continuous line and by a dotted line respectively. We denote the corresponding homotopy classes by $\rho_{ik},\tau_{ik}\in\pi_1(F_{1,n}(Y),\tilde{D})$. Informally ρ_{ik} corresponds to a loop of the i-th point along α_k and τ_{ik} corresponds to a loop of the i-th point along β_k. Let us denote by A_{ij}, $i<j$ the element of $\pi_1(F_{1,n}Y,\tilde{D})$ represented by a closed simple arc in Y^n which leaves fixed b_k for $k\neq i$ and moves b_i along the arc pictured on Figure 2. For every i and every $j>i$ the pictured loop is chosen so as to stay on the left of all arcs used to construct ρ_{ik} and τ_{ik} for $k=1,\dots,g$.



Figure 2: Generators of the pure braid group $\pi_1(F_{1,n},\tilde{D})$

Claim. *The pure braid group* $\pi_1(F_{1,n}Y,\tilde{D})$ *is generated by* ρ_{ik},τ_{ik} *and* A_{ij} *where* $i,j=1,\dots,n,\quad i<j$ *and* $k=1,\dots,g$.

Proof. This is proved by induction on n. When $n=1$ the claim is obvious. Let $n\geq 2$. Consider the fibration $F_{1,n}Y\to F_{1,n-1}Y$ defined by $(y_1,y_2,\dots,y_n)\to(y_2,\dots,y_n)$. One has an exact sequence

$$1\to\pi_1(Y-\{b_0,b_2,\dots,b_n\},b_1)\to\pi_1(F_{1,n}Y,(b_1,\dots,b_n))\to\pi_1(F_{1,n-1}Y,(b_2,\dots,b_n))\to 1$$

since $\pi_2(F_{1,n-1}Y) = 1$ by [FN], Corollary 2.2. The elements $\rho_{ik}, \tau_{ik}, A_{ij}$ with $i \geq 2$ map to the corresponding elements in $\pi_1(F_{1,n-1}Y, (b_2, \dots, b_n))$. The group $\pi_1(Y - \{b_0, b_2, \dots, b_n\}, b_1)$ is freely generated by elements which map into $\rho_{1,k}, \tau_{1,k}, A_{1,j}$ where $j \geq 2, \quad k = 1, \dots, g$. This shows the claim. □

Following [Fu] p.547 and using the notation of 1.2 let us denote by R_i the simply connected region of R enclosed by the arcs ℓ_i and ℓ_{i+1} and the arc of L from b_i to b_{i+1}. For every $i = 1, \dots, n-1$ choose simple arcs $\eta_i : [0,1] \to Y$ going from b_i to b_{i+1} in R_i and $\eta_i' : [0,1] \to Y$ going from b_{i+1} to b_i in R'. Let $s_i : [0,1] \to Y^{(n)} - \Delta$ be the closed arc

$$s_i(t) \ = \ \{b_1, \dots, b_{i-1}, \eta_i(t), \eta_i'(t), b_{i+2}, \dots, b_n\}$$

The homotopy class of s_i is denoted by σ_i. We may consider $\pi_1(F_{1,n}(Y), \tilde{D})$ as embedded in $\pi_1((Y - b_0)^{(n)} - \Delta, D)$. The following relations are easy to verify (cf. [FB] p.249)

$$\begin{aligned} \rho_{i+1,k} \ &= \ \sigma_i \rho_{ik} \sigma_i^{-1}, \quad \tau_{i+1,k} \ = \ \sigma_i \tau_{ik} \sigma_i^{-1} \\ A_{ij} \ &= \ \sigma_i^{-1} \cdots \sigma_{j-2}^{-1} \sigma_{j-1}^2 \sigma_{j-2} \cdots \sigma_i \end{aligned} \tag{7}$$

Proposition 1.6 (Birman). *Let $a, b \in \mathbb{Z}, \quad 1 \leq a, b \leq n$. The braid group $\pi_1((Y - b_0)^{(n)} - \Delta, D)$ is generated by $\sigma_j, \rho_{ak}, \tau_{bk}$ where $1 \leq j \leq n-1, \quad 1 \leq k \leq g$. The corresponding homotopy classes generate $\pi_1(Y^{(n)} - \Delta, D)$ as well.*

Proof. Let us consider the exact sequence Eq.(6). The braids σ_j map to the transpositions $(j \ j+1), \quad j = 1, \dots, n-1$ which generate S_n. So $\sigma_j, \rho_{ik}, \tau_{ik}, A_{ij}$ with $i < j$ generate $\pi_1((Y - b_0)^{(n)} - \Delta, D)$ according to the claim proved in 1.5. The relations Eq.(7) show that the generators may be reduced as stated in the theorem. The last statement follows from the surjection $\pi_1((Y - b_0)^{(n)} - \Delta, D) \twoheadrightarrow \pi_1(Y^{(n)} - \Delta, D)$. □

We described in 1.5 closed arcs in $(Y - b_0)^{(n)} - \Delta$ based at $D = \{b_1, \dots, b_n\}$ whose homotopy classes form a system of generators $\sigma_j, \rho_{ik}, \tau_{ik}$ for $\pi_1((Y - b_0)^{(n)} - \Delta, D)$. Our aim now is for each of these to construct a homotopy of the standard system of closed arcs $\gamma_i^u, \alpha_k^u, \beta_k^u$ as in 1.3. This will permit us to calculate eventually the corresponding braid moves of the Hurwitz systems. The calculation of the braid moves $\sigma_j, \ j = 1, \dots, n-1$ is due to Hurwitz (cf. [Hu] or e.g. [Vo], Theorem 10.3). We define closed arcs $\delta_k : [0,1] \to Y, \ \delta_k(0) = \delta_k(1) = b_0, \ k = 0, 1, \dots, g$ as follows. We let $\delta_0(t) = b_0, \ \forall t \in [0,1]$. We connect the initial vertex of α_1 with the end vertex of β_1^{-1} in the $4g$-polygon of Figure 1 by a simple arc which belongs to the region R' on the right of L (cf. 1.2). This yields δ_1. We connect the initial vertex of α_1 with the end vertex of β_2^{-1} by a simple arc which belongs to the region on the right of L and on the left of δ_1. We denote the corresponding closed arc of Y by δ_2. Continuing in this way we obtain $\delta_1, \delta_2, \dots, \delta_g$ (see Figure 3). Clearly $\delta_k \simeq [\alpha_1, \beta_1] \cdots [\alpha_k, \beta_k]$ in $Y - D$ and $\delta_g \simeq \gamma_1 \cdots \gamma_n$ according to Eq.(2).

Theorem 1.7. *Let Y be a compact, closed Riemann surface of genus $g(Y) \geq 1$. Let $b_0 \in Y$ and let $\gamma_i, \alpha_k, \beta_k$ with $1 \leq i \leq n$, $1 \leq k \leq g$ be a standard system of closed arcs as in 1.2. Let*

$\delta_k, k = 0, 1, \dots, g$ be the closed arcs defined above. For each of the closed arcs in $(Y - b_0)^{(n)} - \Delta$ constructed in 1.5 and representing $\sigma_j, \rho_{ik}, \tau_{ik}$ there is a homotopy $\gamma_i^u, \alpha_k^u, \beta_k^u$, $u \in [0,1]$ of the standard system of closed arcs such that the end system $\gamma'_i, \alpha'_k, \beta'_k$, $1 \le i \le n$, $1 \le k \le g$ is homotopic to:

a. for σ_j where $1 \le j \le n-1$

$$\gamma'_i \simeq \gamma_i \quad \text{for } \forall i \neq j,\, j+1, \quad \alpha'_k \simeq \alpha_k,\ \beta'_k \simeq \beta_k \quad \text{for } \forall k$$
$$\gamma'_j \simeq \gamma_{j+1}, \quad \gamma'_{j+1} \simeq \gamma_{j+1}^{-1}\gamma_j\gamma_{j+1}$$

b. for ρ_{ik} where $1 \le i \le n$, $1 \le k \le g$

$$\gamma'_j \simeq \gamma_j \quad \text{for } \forall j \neq i, \quad \alpha'_\ell \simeq \alpha_\ell \quad \text{for } \forall \ell, \quad \beta'_\ell \simeq \beta_\ell \quad \text{for } \forall \ell \neq k$$
$$\gamma'_i \simeq \mu_{ik}\gamma_i\mu_{ik}^{-1}, \quad \beta'_k \simeq (\zeta_{ik}\gamma_i\zeta_{ik}^{-1})\beta_k, \quad \text{where}$$
$$\mu_{ik} \simeq (\gamma_1 \cdots \gamma_{i-1})^{-1}\delta_{k-1}\alpha_k(\delta_k^{-1}\delta_g)(\gamma_{i+1}\cdots\gamma_n)^{-1}, \quad \zeta_{ik} \simeq (\delta_k^{-1}\delta_g)(\gamma_{i+1}\cdots\gamma_n)^{-1}$$

c. for τ_{ik} where $1 \le i \le n$, $1 \le k \le g$

$$\gamma'_j \simeq \gamma_j \quad \text{for } \forall j \neq i, \quad \alpha'_\ell \simeq \alpha_\ell \quad \text{for } \forall \ell \neq k, \quad \beta'_\ell \simeq \beta_\ell \quad \text{for } \forall \ell$$
$$\gamma'_i \simeq \nu_{ik}\gamma_i\nu_{ik}^{-1}, \quad \alpha'_k \simeq (\xi_{ik}\gamma_i^{-1}\xi_{ik}^{-1})\alpha_k, \quad \text{where}$$
$$\nu_{ik} \simeq \gamma_{i+1}\cdots\gamma_n(\delta_k^{-1}\delta_g)^{-1}\beta_k\delta_{k-1}^{-1}\gamma_1\cdots\gamma_{i-1}, \quad \xi_{ik} \simeq \delta_{k-1}^{-1}\gamma_1\cdots\gamma_{i-1}$$

For each of the inverse closed arcs corresponding to $\sigma_j^{-1}, \rho_{ik}^{-1}, \tau_{ik}^{-1}$ there is a corresponding homotopy of the standard system $\gamma_i, \alpha_k, \beta_k$, $1 \le i \le n$, $1 \le k \le g$ such that the end system $\gamma''_i, \alpha''_k, \beta''_k$, $1 \le i \le n$, $1 \le k \le g$ is homotopic to:

d. for σ_j^{-1} where $1 \le j \le n-1$

$$\gamma''_i \simeq \gamma_i \quad \text{for } \forall i \neq j,\, j+1, \quad \alpha''_k \simeq \alpha_k,\ \beta''_k \simeq \beta_k \quad \text{for } \forall k$$
$$\gamma''_j \simeq \gamma_j\gamma_{j+1}\gamma_j^{-1}, \quad \gamma''_{j+1} \simeq \gamma_j$$

e. for ρ_{ik}^{-1} where $1 \le i \le n$, $1 \le k \le g$

$$\gamma''_j \simeq \gamma_j \quad \text{for } \forall j \neq i, \quad \alpha''_\ell \simeq \alpha_\ell \quad \text{for } \forall \ell, \quad \beta''_\ell \simeq \beta_\ell \quad \text{for } \forall \ell \neq k$$
$$\gamma''_i \simeq \mu_{ik}^{-1}\gamma_i\mu_{ik}, \quad \beta''_k \simeq (\tilde{\zeta}_{ik}^{-1}\gamma_i^{-1}\tilde{\zeta}_{ik})\beta_k, \quad \text{where}$$
$$\mu_{ik} \text{ is as in (b) and } \quad \tilde{\zeta}_{ik} \simeq (\gamma_1\cdots\gamma_{i-1})^{-1}\delta_{k-1}\alpha_k$$

f. for τ_{ik}^{-1} where $1 \le i \le n$, $1 \le k \le g$

$$\gamma''_j \simeq \gamma_j \quad \text{for } \forall j \neq i, \quad \alpha''_\ell \simeq \alpha_\ell \quad \text{for } \forall \ell \neq k, \quad \beta''_\ell \simeq \beta_\ell \quad \text{for } \forall \ell$$
$$\gamma''_i \simeq \nu_{ik}^{-1}\gamma_i\nu_{ik}, \quad \alpha''_k \simeq (\tilde{\xi}_{ik}^{-1}\gamma_i\tilde{\xi}_{ik})\alpha_k, \quad \text{where}$$
$$\nu_{ik} \text{ is as in (c) and } \quad \tilde{\xi}_{ik} \simeq (\gamma_{i+1}\cdots\gamma_n)(\delta_k^{-1}\delta_g)^{-1}\beta_k$$



Figure 3: Homotopy along ρ_{ik}

Proof. (a) Here one moves only γ_j and γ_{j+1}. Clearly $\gamma_j' \simeq \gamma_{j+1}$ and $\gamma_j\gamma_{j+1} \simeq \gamma_j'\gamma_{j+1}'$, so $\gamma_{j+1}' \simeq \gamma_{j+1}^{-1}\gamma_j\gamma_{j+1}$.

(b) The effect of the homotopy of $\gamma_i, \alpha_k, \beta_k$ along ρ_{ik} is pictured on Figure 3. One moves the point b_i along the arc r_{ik} pictured on Figure 2 and together with it deforms the closed arc γ_i. At the moment γ_i^u reaches the side β_k one deforms also β_k in order that the condition γ_i^u and β_k^u have no points in common except b_0 remains valid. None of γ_j for $j \neq i$, or α_ℓ for $\forall \ell$, or β_ℓ for $\forall \ell \neq k$ changes in this homotopy. The effect of cutting Y along the closed arcs $\alpha_1, \beta_1, \ldots, \alpha_k, \beta_k', \ldots, \alpha_g, \beta_g$ is the same as to cut a region containing b_i from the original $4g$-polygon and glue it along the side β_i as described in Figure 3 (NW) (NW=Northwest). We wish to express γ_i' in terms of the standard system $\gamma_j, \alpha_\ell, \beta_\ell$, $1 \leq j \leq n, 1 \leq \ell \leq g$. The closed arc γ_i' is pictured in the original $4g$-polygon on Figure 3 (NE). It is clear that $\gamma_i' \simeq \eta_{ik}\gamma_i\eta_{ik}^{-1}$ where η_{ik} is the closed arc pictured on Figure 3 (SW). That η_{ik} is homotopic to $(\gamma_1 \cdots \gamma_{i-1})^{-1}(\delta_{k-1}\alpha_k)\delta_k^{-1}(\gamma_1 \cdots \gamma_{i-1})$ is evident from Figure 3 (SE). Furthermore $\eta_{ik}\gamma_i\eta_{ik}^{-1} \simeq (\eta_{ik}\gamma_i)\gamma_i(\eta_{ik}\gamma_i)^{-1}$. Using the relation $\gamma_1 \cdots \gamma_n \simeq \delta_g$ we obtain $\eta_{ik}\gamma_i \simeq (\gamma_1 \cdots \gamma_{i-1})^{-1}\delta_{k-1}\alpha_k(\delta_k^{-1}\delta_g)(\gamma_{i+1} \cdots \gamma_n)^{-1}$. This proves the formula for γ_i' of Part (b). The calculation of β_k' is similar. The closed arc β_k' is pictured in the original $4g$-polygon on Figure 4 (NW). It is homotopic to $\omega_{ik} \cdot \beta_k$ where ω_{ik}



Figure 4:

is the closed arc based at b_0 pictured on Figure 4 (NE). We then consider $\delta_k \omega_{ik} \delta_k^{-1}$ (see Figure 4 (SW)). The latter is homotopic to $(\gamma_1 \cdots \gamma_{i-1})\gamma_i(\gamma_1 \cdots \gamma_{i-1})^{-1}$. We thus obtain $\omega_{ik} \simeq (\delta_k^{-1}\gamma_1 \cdots \gamma_{i-1})\gamma_i(\gamma_1 \cdots \gamma_1)^{-1}\delta_k$. We then have

$$\begin{aligned}\omega_{ik} &\simeq \delta_k^{-1}(\gamma_1 \cdots \gamma_{i-1}\gamma_i)\gamma_i(\gamma_1 \cdots \gamma_{i-1}\gamma_i)^{-1}\delta_k \\ &\simeq [\delta_k^{-1}\delta_g(\gamma_{i+1} \cdots \gamma_n)^{-1}]\gamma_i[\gamma_{i+1} \cdots \gamma_n \delta_g^{-1}\delta_k]\end{aligned}$$

since $\gamma_1 \cdots \gamma_n \simeq \delta_g$. This proves the second formula of Part (b).

(c) The arguments here are very similar to those of Part (b). One deforms γ_i and α_k with b_i^u moving along the arc t_{ik} (see Figure 2). When b_i^u returns to b_i one obtains closed arcs γ_i' and α_k' for which $\gamma_i' \simeq \theta_{ik}\gamma_i\theta_{ik}^{-1}$, $\alpha_k' = \varepsilon_{ik}\alpha_k$ where θ_{ik} and ε_{ik} are represented by arcs in the original $4g$-polygon pictured on Figure 5. We calculate θ_{ik} in the following way. We consider it as a product of four arcs according to the picture. We deform θ_{ik} in such a way that the first arc becomes a closed simple arc encircling $\{b_1, \ldots, b_i\}$ in clockwise direction, then the second arc goes from the initial point of α_1 to the end point of α_k^{-1}, the third arc goes from the initial point of α_k to the initial point of α_1 and the fourth arc equals the first one with the opposite orientation. We have accordingly

$$\theta_{ik} \simeq (\gamma_1 \cdots \gamma_i)^{-1}(\delta_k\beta_k)\delta_{k-1}^{-1}(\gamma_1 \cdots \gamma_i)$$



Figure 5:

Conjugating γ_i by θ_{ik} we may cancel the last factor γ_i from θ_{ik} and replace $(\gamma_1 \cdots \gamma_i)^{-1}$ by $(\gamma_{i+1} \cdots \gamma_n)\delta_g^{-1}$. We thus obtain $\gamma_i' = \nu_{ik}\gamma_i\nu_{ik}^{-1}$ where

$$\nu_{ik} \ \simeq \ (\gamma_{i+1} \cdots \gamma_n)(\delta_k^{-1}\delta_g)^{-1}\beta_k\delta_{k-1}^{-1}(\gamma_1 \cdots \gamma_{i-1})$$

Finally, $\delta_{k-1}\varepsilon_{ik}\delta_{k-1}^{-1} \simeq (\gamma_1 \cdots \gamma_{i-1})\gamma_i^{-1}(\gamma_1 \cdots \gamma_{i-1})^{-1}$. This proves the last formula of Part (c).

In order to obtain the formulas of (d), (e) and (f) from those of (a), (b) and (c) we notice that if we apply in each case to $(\gamma_1'', \ldots, \gamma_n''; \alpha_1'', \ldots, \beta_g'')$ the braid moves $\sigma_j, \rho_{ik}, \tau_{ik}$ respectively we obtain $(\gamma_1, \ldots, \gamma_n; \alpha_1, \ldots, \beta_g)$. For example, in order to verify (d) we have $\gamma_j \simeq \gamma_{j+1}''$, $\gamma_{j+1} \simeq (\gamma_{j+1}'')^{-1}\gamma_j''\gamma_{j+1}''$. Therefore $\gamma_j'' \simeq \gamma_j\gamma_{j+1}\gamma_j^{-1}$. In Case (e) applying ρ_{ik} to $(\gamma_1'', \ldots, \beta_g'')$ we obtain $\gamma_j \simeq \gamma_j''$ for $\forall j \neq i$, $\alpha_\ell \simeq \alpha_\ell''$ for $\forall \ell$, $\beta_\ell \simeq \beta_\ell''$ for $\forall \ell \neq k$. Furthermore $\gamma_i \simeq \mu_{ik}''\gamma_i''(\mu_{ik}'')^{-1}$ where μ_{ik}'' is expressed by $\gamma_j'', \alpha_\ell'', \beta_\ell''$ as in (b). Since neither γ_i'' nor β_k'' enter in this expression we may replace $\gamma_j'', \alpha_\ell'', \beta_\ell''$ by $\gamma_j, \alpha_\ell, \beta_\ell$ and we obtain $\mu_{ik}'' = \mu_{ik}$. Thus $\gamma_i'' \simeq \mu_{ik}^{-1}\gamma_i\mu_{ik}$. Similarly we have $\beta_k = (\zeta_{ik}\gamma_i''\zeta_k^{-1})\beta_k''$ where ζ_{ik} is as in (b). Replacing γ_i'' by $\mu_{ik}^{-1}\gamma_i\mu_{ik}$ and canceling we obtain $\beta_k'' = (\tilde{\zeta}_{ik}^{-1}\gamma_i^{-1}\tilde{\zeta}_{ik})\beta_k$ where $\tilde{\zeta}_{ik}$ is as in (e). In a similar manner one deduces (f) from (c). ☐

Recall from 1.1 and 1.2 that given $D \in Y^{(n)} - \Delta$, $b_0 \in Y - D$ and fixing a standard system of closed arcs $\gamma_1, \ldots, \gamma_n; \alpha_1, \beta_1, \ldots, \alpha_g, \beta_g$ there is a bijective correspondence between the fiber of $\mathcal{H}_d^{(n)}(Y, b_0) \to (Y - b_0)^{(n)} - \Delta$ over D and the set of Hurwitz systems $(t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$. Similarly there is a bijective correspondence between the fiber of $\mathcal{U}(b_0) \to (Y - b_0)^{(n)} - \Delta$, where $\mathcal{U}(b_0) \subset \mathcal{H}_d^{(n)}(Y)$, and the set of equivalence classes of Hurwitz systems $[t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g]$. In the next theorem we calculate the monodromy action of $\pi_1((Y - b_0)^{(n)} - \Delta, D)$ on these fibers. According to Proposition 1.6 it suffices to determine the braid moves which correspond to the generators $\sigma_j, \rho_{ik}, \tau_{ik}$. We denote the corresponding braid moves of the first type (cf. Definition 2) by $\sigma_j', \rho_{ik}', \tau_{ik}'$ and we denote the corresponding braid moves of the second type (inverse to those of the first type) by $\sigma_j'', \rho_{ik}'', \tau_{ik}''$.

Theorem 1.8. *Let* $(t_1,\dots,t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$, $\lambda_{g+k}=\mu_k$, *be a Hurwitz system. Let* $u_k=[\lambda_1,\mu_1]\cdots[\lambda_k,\mu_k]$ *for* $k=1,\dots,g$ *and let* $u_0=1$. *The following formulas hold for the braid moves* $(t_1,\dots,t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)\mapsto(t'_1,\dots,t'_n;\lambda'_1,\mu'_1,\dots,\lambda'_g,\mu'_g)$ *of the first type.*

a. *For* σ'_j *where* $1\le j\le n-1$

$$t'_i=t_i \quad \text{for } \forall i\neq j,\ j+1,\quad \lambda'_\ell=\lambda_\ell,\quad \mu'_\ell=\mu_\ell \quad \text{for } \forall\ell$$
$$(t_j,t_{j+1})\ \mapsto\ (t'_j,t'_{j+1})\ =\ (t_jt_{j+1}t_j^{-1}\,,\ t_j). \tag{8}$$

b. *For* ρ'_{ik} *where* $1\le i\le n,\ 1\le k\le g$

$$t'_j=t_j \quad \text{for } \forall j\neq i,\quad \lambda'_\ell=\lambda_\ell \quad \text{for } \forall\ell,\quad \mu'_\ell=\mu_\ell \quad \text{for } \forall\ell\neq k$$
$$(t_i,\mu_k)\ \mapsto\ (t'_i,\mu'_k)\ =\ (a_1^{-1}t_ia_1,(b_1^{-1}t_i^{-1}b_1)\mu_k) \quad \text{where}$$
$$a_1=(t_1\cdots t_{i-1})^{-1}u_{k-1}\lambda_k(u_k^{-1}u_g)(t_{i+1}\cdots t_n)^{-1},\quad b_1=(t_1\cdots t_{i-1})^{-1}u_{k-1}\lambda_k.$$

c. *For* τ'_{ik} *where* $1\le i\le n,\ 1\le k\le g$

$$t'_j=t_j \quad \text{for } \forall j\neq i,\quad \lambda'_\ell=\lambda_\ell \quad \text{for } \forall\ell\neq k,\quad \mu'_\ell=\mu_\ell \quad \text{for } \forall\ell$$
$$(t_i,\lambda_k)\ \mapsto\ (t'_i,\lambda'_k)\ =\ (c_1^{-1}t_ic_1,(d_1^{-1}t_id_1)\lambda_k) \quad \text{where}$$
$$c_1=t_{i+1}\cdots t_n(u_k^{-1}u_g)^{-1}\mu_k(u_{k-1})^{-1}t_1\cdots t_{i-1},\quad d_1=t_{i+1}\cdots t_n(u_k^{-1}u_g)^{-1}\mu_k.$$

The following formulas hold for the braid moves $(t_1,\dots,t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)\mapsto(t''_1,\dots,t''_n;\lambda''_1,\mu''_1,\dots,\lambda''_g,\mu''_g)$ *of the second type.*

d. *For* σ''_j *where* $1\le j\le n-1$

$$t''_i=t_i \quad \text{for } \forall i\neq j,\ j+1,\quad \lambda''_\ell=\lambda_\ell,\quad \mu''_\ell=\mu_\ell \quad \text{for } \forall\ell$$
$$(t_j,t_{j+1})\ \mapsto\ (t''_j,t''_{j+1})\ =\ (t_{j+1}\,,\ t_{j+1}^{-1}t_jt_{j+1}). \tag{9}$$

e. *For* ρ''_{ik} *where* $1\le i\le n,\ 1\le k\le g$

$$t''_j=t_j \quad \text{for } \forall j\neq i,\quad \lambda''_\ell=\lambda_\ell \quad \text{for } \forall\ell,\quad \mu''_\ell=\mu_\ell \quad \text{for } \forall\ell\neq k$$
$$(t_i,\mu_k)\ \mapsto\ (t''_i,\mu''_k)\ =\ (a_2^{-1}t_ia_2,(b_2^{-1}t_ib_2)\mu_k) \quad \text{where}$$
$$a_2=t_{i+1}\cdots t_n(u_k^{-1}u_g)^{-1}\lambda_k^{-1}(u_{k-1})^{-1}t_1\cdots t_{i-1},\quad b_2=t_{i+1}\cdots t_n(u_k^{-1}u_g)^{-1}.$$

f. *For* τ''_{ik} *where* $1\le i\le n,\ 1\le k\le g$

$$t''_j=t_j \quad \text{for } \forall j\neq i,\quad \lambda''_\ell=\lambda_\ell \quad \text{for } \forall\ell\neq k,\quad \mu''_\ell=\mu_\ell \quad \text{for } \forall\ell$$
$$(t_i,\lambda_k)\ \mapsto\ (t''_i,\lambda''_k)\ =\ (c_2^{-1}t_ic_2,(d_2^{-1}t_i^{-1}d_2)\lambda_k) \quad \text{where}$$
$$c_2=(t_1\cdots t_{i-1})^{-1}u_{k-1}\mu_k^{-1}(u_k^{-1}u_g)(t_{i+1}\cdots t_n)^{-1},\quad d_2=(t_1\cdots t_{i-1})^{-1}u_{k-1}.$$

Proof. Formulas (d), (e) and (f) are obtained from formulas (a), (b) and (c) of Theorem 1.7 respectively applying the homomorphism $m:\pi_1(Y-D,b_0)\to S_d$ and equalities Eq.(4) and Eq.(5). By Lemma 1.4 the braid moves of the first type are inverse to the braid moves of the second type. We may thus obtain formulas (a), (b) and (c) from formulas (d), (e) and (f) of Theorem 1.7 respectively applying the homomorphism m. ☐

Corollary 1.9. *Using the notation of Theorem 1.8 the following formulas hold.*

i. For ρ'_{ik} *:*

$$\begin{aligned}
&\rho'_{nk} : t_n \mapsto t'_n = a_1^{-1} t_n a_1, && \textit{where } a_1 = (u_k^{-1}u_g)^{-1}[\mu_k, \lambda_k]\lambda_k(u_k^{-1}u_g)\\
&\rho'_{1k} : \mu_k \mapsto \mu'_k = (b_1^{-1}t_1^{-1}b_1)\mu_k, && \textit{where } b_1 = u_{k-1}\lambda_k\\
&\rho'_{nk} : \mu_k \mapsto \mu'_k = (b_1^{-1}t_n^{-1}b_1)\mu_k, && \textit{where } b_1 = (u_k^{-1}u_g)^{-1}[\mu_k, \lambda_k]\lambda_k.
\end{aligned}$$

ii. For τ'_{ik} *:*

$$\begin{aligned}
&\tau'_{1k} : t_1 \mapsto t'_1 = c_1^{-1} t_1 c_1, && \textit{where } c_1 = u_{k-1}[\lambda_k, \mu_k]\mu_k u_{k-1}^{-1}\\
&\tau'_{1k} : \lambda_k \mapsto \lambda'_k = (d_1^{-1}t_1 d_1)\lambda_k, && \textit{where } d_1 = u_{k-1}[\lambda_k, \mu_k]\mu_k\\
&\tau'_{nk} : \lambda_k \mapsto \lambda'_k = (d_1^{-1}t_n d_1)\lambda_k, && \textit{where } d_1 = (u_k^{-1}u_g)^{-1}\mu_k.
\end{aligned}$$

iii. For ρ''_{ik} *:*

$$\begin{aligned}
&\rho''_{1k} : t_1 \mapsto t''_1 = a_2^{-1} t_1 a_2, && \textit{where } a_2 = u_{k-1}[\lambda_k, \mu_k]\lambda_k^{-1}u_{k-1}^{-1}\\
&\rho''_{1k} : \mu_k \mapsto \mu''_k = (u_k^{-1}t_1 u_k)\mu_k, &&\\
&\rho''_{nk} : \mu_k \mapsto \mu''_k = (b_1^{-1}t_n b_1)\mu_k, && \textit{where } b_1 = (u_k^{-1}u_g)^{-1}.
\end{aligned}$$

iv. For τ''_{ik} *:*

$$\begin{aligned}
&\tau''_{nk} : t_n \mapsto t''_n = c_2^{-1} t_n c_2, && \textit{where } c_2 = (u_k^{-1}u_g)^{-1}[\mu_k, \lambda_k]\mu_k^{-1}(u_k^{-1}u_g)\\
&\tau''_{1k} : \lambda_k \mapsto \lambda''_k = (u_{k-1}^{-1}t_1^{-1}u_{k-1})\lambda_k, &&\\
&\tau''_{nk} : \lambda_k \mapsto \lambda''_k = (d_2^{-1}t_n^{-1}d_2)\lambda_k, && \textit{where } d_2 = (u_k^{-1}u_g)^{-1}[\mu_k, \lambda_k].
\end{aligned}$$

In particular

$$\rho''_{ng} \;:\; \mu_g \;\mapsto\; t_n\mu_g, \qquad \tau''_{11} \;:\; \lambda_1 \;\mapsto\; t_1^{-1}\lambda_1. \tag{10}$$

Proof. Let us prove the first formula of Part (ii). The other formulas can be either proved similarly or are restatements of particular cases of Theorem 1.8. We have $\tau'_{1k} : t_1 \mapsto t'_1 = c_1^{-1}t_1c_1$ where $c_1 = t_2\cdots t_n(u_k^{-1}u_g)^{-1}\mu_k u_{k-1}^{-1}$. Since $t_1\cdots t_n = u_g$ it holds $(t_2\cdots t_n)^{-1}t_1(t_2\cdots t_n) = u_g^{-1}t_1u_g$. Hence

$$\begin{aligned}
t'_1 \;&=\; (u_{k-1}\mu_k^{-1}u_k^{-1})t_1(\cdots)^{-1} \;=\; (u_{k-1}\mu_k^{-1}[\lambda_k, \mu_k]^{-1}u_{k-1}^{-1})t_1(\cdots)^{-1}\\
&=\; \left(u_{k-1}[\lambda_k, \mu_k]\mu_k u_{k-1}^{-1}\right)^{-1} t_1 \left(u_{k-1}[\lambda_k, \mu_k]\mu_k u_{k-1}^{-1}\right)
\end{aligned}$$

□

Summing up the discussion made so far in this section we obtain the following result.

Theorem 1.10. *Fix* $d \geq 2$, $n > 0$, $g \geq 0$. *Let us consider the set of all Hurwitz systems* $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ *(cf. Definition 1). Let* F *be the free group generated by the symbols* $\sigma_j, \rho_{ik}, \tau_{ik}$ *where* $1 \leq j \leq n-1$, $1 \leq i \leq n$, $1 \leq k \leq g$. *Let us consider the right action of* F *on the set of Hurwitz systems defined by the formulas for* $\sigma'_j, \rho'_{ik}, \tau'_{ik}$ *of Theorem 1.8 (a)–(c) or alternatively let us consider the left action of* F *on the set of Hurwitz systems defined by the formulas for* $\sigma''_j, \rho''_{ik}, \tau''_{ik}$ *of Theorem 1.8 (d)–(f). Then the connected components of* $\mathcal{H}^{(n)}_d(Y, b_0)$ *(cf. 1.1) correspond bijectively to the orbits of either action of* F *and the connected components of* $\mathcal{H}^{(n)}_d(Y)$ *correspond bijectively to the orbits of either associated action of* F *on the set of equivalence classes of Hurwitz systems modulo inner automorphisms of* S_d.

Proof. $\mathcal{H}^{(n)}_d(Y, b_0) \to (Y - b_0)^{(n)} - \Delta$ is a topological covering map where a Hausdorff topology on $\mathcal{H}^{(n)}_d(Y, b_0)$ is defined as in [Fu] pp.545,546. According to the definition of product of arcs Eq.(1) the monodromy action of the fundamental group $\pi_1((Y - b_0)^{(n)} - \Delta, D)$ on the fiber over D is a right action. The identification of this fiber with the set of Hurwitz systems, Proposition 1.6 and the calculation of the braid moves of the first type $\sigma'_j, \rho'_{ik}, \tau'_{ik}$ in Theorem 1.8 yield the statement about the orbits of the right action of F.

Consider the associated left action $gx = xg^{-1}$. The orbits are the same and according to Lemma 1.4 the braid moves of the second type $\sigma''_j, \rho''_{ik}, \tau''_{ik}$ are inverse to those of the first type $\sigma'_j, \rho'_{ik}, \tau'_{ik}$ respectively. This shows the statement about the orbits of the left action of F.

Let $b_0 \in Y$. Since $\mathcal{U}(b_0) \subset \mathcal{H}^{(n)}_d(Y)$ (cf. 1.1) is a Zariski dense open subset, the connected components of $\mathcal{U}(b_0)$ correspond bijectively to those of $\mathcal{H}^{(n)}_d(Y)$. This shows the last claim of the theorem. □

2 The main lemma

In the previous section we considered Hurwitz systems $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ where $t_i, \lambda_k, \mu_k \in S_d$ in connection with the problem of determining the connected components of the Hurwitz spaces. In this section we replace S_d by an arbitrary (possibly infinite) group G.

Definition 3. Let G be an arbitrary group. We call $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ a Hurwitz system with values in G if $t_i, \lambda_k, \mu_k \in G$, $t_i \neq 1$ for $\forall i$ and $t_1 \cdots t_n = [\lambda_1, \mu_1] \cdots [\lambda_g, \mu_g]$.

The formulas of Theorem 1.8 make sense for an arbitrary group G. That $\sigma''_j = (\sigma'_j)^{-1}$, $\rho''_{ik} = (\rho'_{ik})^{-1}$ and $\tau''_{ik} = (\tau'_{ik})^{-1}$ is evident from Theorem 1.7.

Definition 4. We call two Hurwitz systems with values in G braid-equivalent if one is obtained from the other by a finite sequence of braid moves $\sigma'_j, \rho'_{ik}, \tau'_{ik}, \sigma''_j, \rho''_{ik}, \tau''_{ik}$ where $1 \leq j \leq n-1$, $1 \leq i \leq n$, $1 \leq k \leq g$. We denote the braid equivalence by $\sim$.

Main Lemma 2.1. *Let G be an arbitrary group and let $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ be a Hurwitz system with values in G. Suppose that $t_i t_{i+1} = 1$. Let H be the subgroup of G generated by $\{t_1, \dots, t_{i-1}, t_{i+2}, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g\}$. Then for every $h \in H$ the given Hurwitz system is braid-equivalent to $(t_1, \dots, t_{i-1}, t_i^h, t_{i+1}^h, t_{i+2}, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$*

Proof. Let us fix $t_1, \dots, t_{i-1}, t_{i+2}, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g$. Let $H_1 \subseteq H$ be the subset consisting of elements h such that the statement of the lemma holds for an arbitrary pair $(t_i, t_{i+1}) = (\tau, \tau^{-1})$.

Step 1. *We claim H_1 is a subgroup of H.* Let $h \in H_1$ and let $t = h t_i h^{-1}$. Then $t_i = t^h$, so by assumption $(\dots, t^h, (t^{-1})^h, \dots)$ can be obtained from $(\dots, t, t^{-1}, \dots)$ by a sequence of braid moves. Then one can obtain $(\dots, t, t^{-1}, \dots)$ from $(\dots, t_i, t_i^{-1}, \dots)$ by the inverse sequence of braid moves. Thus $h^{-1} \in H_1$. If $h_1, h_2 \in H_1$, then $(\dots, t_i, t_i^{-1}, \dots) \sim (\dots, t_i^{h_1}, (t_i^{-1})^{h_1}, \dots) \sim (\dots, t_i^{h_1 h_2}, (t_i^{-1})^{h_1 h_2}, \dots)$, so $h_1 h_2 \in H_1$.

Step 2. *For every $\ell \neq i, i+1$ the element t_ℓ belongs to H_1.* Applying a sequence of braid moves σ_j', σ_j'' we can move the adjacent pair (t_i, t_{i+1}) wherever we want among the first n elements of the Hurwitz system without changing the other elements. So move (t_i, t_{i+1}) to the left side of t_ℓ. Then we have $(t_i, t_{i+1}, t_\ell) \sim (t_i, t_\ell, t_\ell^{-1} t_{i+1} t_\ell) \sim (t_\ell, t_\ell^{-1} t_i t_\ell, t_\ell^{-1} t_{i+1} t_\ell)$. We then move the pair (t_i^h, t_{i+1}^h) with $h = t_\ell$ back to the initial position.

Step 3. *For every $k = 1, \dots, g$ the element $h = u_{k-1} \lambda_k u_k^{-1}$ belongs to H_1.* First suppose that $i = 1$. In this case $t_2 = t_1^{-1}$. Let us perform a braid move ρ_{1k}'. One obtains $(t_1, t_1^{-1}, \dots, \lambda_k, \mu_k, \dots) \sim (t_1', t_1^{-1}, \dots, \lambda_k, \mu_k', \dots)$ where $t_1' = a_1^{-1} t_1 a_1$ with $a_1 = u_{k-1} \lambda_k (u_k^{-1} u_g)(t_2 \cdots t_n)^{-1}$ and $\mu_k' = (b_1^{-1} t_1^{-1} b_1) \mu_k$ with $b_1 = u_{k-1} \lambda_k$. We have $t_1 \cdots t_n = u_g$, so $a_1 = u_{k-1} \lambda_k u_k^{-1} t_1$. Let us move $t_2 = t_1^{-1}$ to the first place using σ_1''. One obtains $(t_1^{-1}, t_1 t_1' t_1^{-1}, \dots, \lambda_k, \mu_k', \dots)$. Here $t_1 t_1' t_1^{-1} = t_1^h$ where $h = u_{k-1} \lambda_k u_k^{-1}$. Let us perform again ρ_{1k}'. The elements u_{k-1}, λ_k have not changed with respect to the original Hurwitz system, so $\mu_k' \mapsto [(u_{k-1}\lambda_k)^{-1} (t_1^{-1})^{-1} (u_{k-1} \lambda_k)] \mu_k' = \mu_k$. Moving the second element t_1^h to the first place by σ_1'' we obtain $(t_1^h, \tilde{t}_2, t_3, \dots, \lambda_k, \mu_k, \dots)$. Since $t_1 t_2 t_3 \cdots t_n = u_g = t_1^h \tilde{t}_2 t_3 \cdots t_n$ we must have $\tilde{t}_2 = (t_1^{-1})^h$. This proves $(t_1, t_1^{-1}, t_3, \dots) \sim (t_1^h, (t_1^{-1})^h, t_3, \dots)$ with $h = u_{k-1} \lambda_k u_k^{-1}$. One extends this braid equivalence to every adjacent pair (t_i, t_{i+1}) with $t_i t_{i+1} = 1$ by moving first the pair to the front, applying the braid equivalence we have just proved and moving the obtained pair back to the original position.

Step 4. *For every $k = 1, \dots, g$ the element $h = u_{k-1} \mu_k^{-1} u_k^{-1}$ belongs to H_1.* The proof is the same as that of Step 3. One uses the braid move τ_{1k}'' instead of ρ_{1k}'.

Step 5. By the preceding steps it remains to verify that λ_k, μ_k for $\forall k$ belong to the subgroup $H_2 \subseteq H_1$ generated by $t_j, u_{k-1} \lambda_k u_k^{-1}, u_{k-1} \mu_k^{-1} u_k^{-1}$ where $j = 1, \dots, i-1, i+2, \dots, n$ and $k = 1, \dots, g$. We prove this by induction on k. We have $\lambda_1 u_1^{-1} \in H_2$, $u_1 \mu_1 = (\mu_1^{-1} u_1^{-1})^{-1} \in H_2$, so $\lambda_1 \mu_1 \in H_2$. Furthermore $H_2 \ni u_1 \mu_1 = [\lambda_1, \mu_1] \mu_1 = \lambda_1 \mu_1 \lambda_1^{-1}$. So $\lambda_1^{\pm 1} \in H_2$ and $\mu_1^{\pm 1} \in H_2$. Let $k \geq 2$. Suppose, by inductive assumption, that $\lambda_1, \mu_1, \dots, \lambda_{k-1}, \mu_{k-1}$ belong to H_2. Then $u_{k-1} = \prod_{\ell=1}^{k-1} [\lambda_\ell, \mu_\ell] \in H_2$. We have $u_{k-1} \lambda_k u_k^{-1} \in H_2$, $u_k \mu_k u_{k-1}^{-1} = (u_{k-1} \mu_k^{-1} u_k^{-1})^{-1} \in H_2$, so $u_{k-1} \lambda_k \mu_k u_{k-1}^{-1} \in H_2$ and therefore $\lambda_k \mu_k \in H_2$. Furthermore $H_2 \ni u_k \mu_k u_{k-1}^{-1} = u_{k-1} [\lambda_k, \mu_k] \mu_k u_{k-1}^{-1}$. Therefore $\lambda_k \mu_k \lambda_k^{-1} \in H_2$, so $\lambda_k^{\pm 1} \in H_2$ and $\mu_k^{\pm 1} \in H_2$.

The lemma is proved. $\square$

3 The case $n \geq 2d-2$

So far we have not made any restrictions on the type of the covering $\pi : X \to Y$. We shall be further occupied mainly with simply branched coverings with connected X. These coverings correspond to Hurwitz systems $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ with local monodromies $t_i = m(\gamma_i)$, $i = 1, \dots, n$ equal to transpositions and with transitive monodromy group $G = \langle t_1, \dots, t_n, \lambda_1, \mu_1, \dots, \lambda_g, \mu_g \rangle$. We call such coverings *simple coverings.* By Hurwitz' formula $n \equiv 0 (mod\, 2)$ for a simple covering.

Definition 5. Let Y be a smooth projective curve. We denote by $\mathcal{H}_{d,n}(Y, b_0)$ and $\mathcal{H}_{d,n}(Y)$ the Hurwitz spaces which parametrize equivalence classes $[X \to Y,\ \phi]$ and $[X \to Y]$ respectively of simple coverings of degree d branched in n points (cf. 1.1).

As a first application of Theorem 1.10 we give a proof, using braid moves, of the following well-known fact.

Proposition 3.1. *Let* $n > 0$, $n \equiv 0(mod\, 2)$. *The Hurwitz spaces* $\mathcal{H}_{2,n}(Y, b_0)$ *and* $\mathcal{H}_{2,n}(Y)$ *which parametrize double coverings of* Y *branched in* n *points are both irreducible.*

Proof. Since $\mathcal{H}_{2,n}(Y, b_0)$ and $\mathcal{H}_{2,n}(Y)$ are smooth it suffices to prove they are connected. Let us first consider $\mathcal{H}_{2,n}(Y, b_0)$. By Theorem 1.10 it suffices to prove that every Hurwitz system $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ with $t_i, \lambda_k, \mu_k \in S_2$ is braid-equivalent to $((12), \dots, (12); 1, 1, \dots, 1, 1)$. According to Theorem 1.8 one has $\rho'_{ik} : (t_i, \mu_k) \mapsto (t_i, t_i\mu_k)$, $\tau'_{ik} : (t_i, \lambda_k) \mapsto (t_i, t_i\lambda_k)$ and the same formulas hold for ρ''_{ik} and τ''_{ik}. Hence whenever some $\lambda_k \neq 1$ or $\mu_k \neq 1$ we can perform a braid move τ'_{1k} or ρ'_{1k} respectively or both in order to obtain a new Hurwitz system with $\lambda'_k = \mu'_k = 1$. The connectedness of $\mathcal{H}_{2,n}(Y, b_0)$ implies the connectedness of the Zariski dense open subset $\mathcal{U}(b_0) \subset \mathcal{H}_{2,n}(Y)$. Therefore $\mathcal{H}_{2,n}(Y)$ is connected as well. $\square$

Given an ordered n-tuple of permutations $\underline{t} = (t_1, \dots, t_n)$ whose product $t_1 \cdots t_n = s \in S_d$, performing an elementary move σ'_j or σ''_j one obtains a new ordered n-tuple $(t'_1, \dots, t'_n)$ which has the same product $t'_1 \cdots t'_n = s$. We shall also call ordered n-tuples sequences.

Definition 6. Two ordered n-tuples (or sequences) of permutations $\underline{t} = (t_1, \dots, t_n)$ and $\underline{t}' = (t'_1, \dots, t'_n)$ are called braid-equivalent if $\underline{t}'$ is obtained from $\underline{t}$ by a finite sequence of braid moves of type σ'_j or σ''_j. We write $\underline{t}' \sim \underline{t}$.

Notice the difference between braid equivalence of sequences and that of Hurwitz systems (cf. Definition 4). The latter includes also braid moves of the types $\rho'_{ik}, \rho''_{ik}, \tau'_{ik}, \tau''_{ik}$.

3.2. We recall some results proved in [Mo]. Given a permutation $s \in S_d$ one considers an ordered n-tuple of transpositions $\underline{t} = (t_1, \dots, t_n)$ such that $t_1 \cdots t_n = s$. Let $\Sigma_1, \Sigma_2, \dots, \Sigma_m$ be the domains of transitivity of the group $G = \langle t_1, \dots, t_n \rangle$. Then $G = S(\Sigma_1) \times \cdots \times S(\Sigma_m)$. For reader's convenience we give a proof of the following important lemma due to Mochizuki (cf. [Mo] Lemma 2.4).

Lemma 3.3. *Let* $\underline{t} = (t_1, \ldots, t_n)$, $t_1 \cdots t_n = s$ *be as above. Let* $a, b \in \Sigma_i$, $a < b$. *Then* $\underline{t} \sim \underline{t}' = ((ab), t'_2, \ldots)$ *and* $\underline{t} \sim \underline{t}'' = (\ldots, t''_{n-1}, (ab))$.

Proof. If (t_i, t_{i+1}) is a pair such that $t_i \in S(\Sigma_k)$, $t_{i+1} \in S(\Sigma_\ell)$ with $k \neq \ell$, then $(t_i, t_{i+1}) \sim (t_{i+1}, t_{i+1}^{-1} t_i t_{i+1}) = (t_{i+1}, t_i)$. This shows that performing a sequence of braid moves (of type σ'_j, σ''_j) one may replace $\underline{t}$ by a concatenation of sequences $T_1 T_2 \ldots T_m$ where T_i is formed by all transpositions of $\underline{t}$ which belong to $S(\Sigma_j)$ ordered in the way they appear in $\underline{t}$ (if $\# \Sigma_j = 1$ we let $T_j = \emptyset$). Furthermore we may move any T_i to the first place or to the last place. This shows that it suffices to prove the lemma in the particular case when $\langle t_1, \ldots, t_n \rangle$ is a transitive group. Let $b = a^\tau$ where $\tau = t_{j_1} t_{j_2} \cdots t_{j_\ell}$. One may vary the sequence within the set of sequences braid-equivalent to $\underline{t}$. Given a sequence one may vary τ so that $b = a^\tau$ and finally given a sequence and a τ one may vary the factorization of τ. Let $\underline{t}'$, τ' with the property $a^{\tau'} = b$ and a factorization $\tau' = t'_{i_1} \cdots t'_{i_r}$ be chosen so that r is minimal possible. If $r = 1$ one has that (ab) is one of the transpositions of $\underline{t}'$. Moving (ab) to the front by subsequent elementary moves σ''_j one obtains a braid-equivalent sequence of the type $((ab), \ldots)$. Moving (ab) to the end by subsequent elementary moves σ'_j one obtains $(\ldots, (ab))$ as required in the lemma. Suppose that $r \geq 2$. Let $x_1 = a$, $x_2 = a^{t'_{i_1}}, \ldots, x_{k+1} = a^{t'_{i_1} \cdots t'_{i_k}}, \ldots, x_{r+1} = b$. The minimality of r implies that $x_i \neq x_j$ for $i \neq j$, so $t'_{i_k} = (x_k \, x_{k+1})$ for $k = 1, \ldots, r$ are r different transpositions of the sequence $\underline{t}'$ and every x_i for $2 \leq i \leq r$ enters in at most two transpositions of the set $\{t'_{i_k}\}$. Applying several elementary moves one places $(a x_2)$ adjacent to $(x_2 x_3)$. One obtains a sequence either of the type $\ldots, (a \, x_2), (x_2 x_3), \ldots$ or of the type $\ldots, (x_2 x_3), (a \, x_2), \ldots$ in which none of t'_{i_k}, $k = 1, \ldots, r$ has been changed. In the first case one has $((a \, x_2), (x_2 x_3)) \sim ((x_2 x_3), (a \, x_3))$ and in the second case $((x_2 x_3), (a \, x_2)) \sim ((a \, x_3), (x_2 x_3))$. In both cases one obtains a braid-equivalent sequence for which the sequence $a = x_1, x_2, x_3, \ldots, x_{r+1} = b$ is replaced by $a = x_1, x_3, \ldots, x_{r+1} = b$. This contradicts the minimality of r. $\square$

3.4. Let $(t_1, \ldots, t_n)$ be an n-tuple of transpositions and let $t_1 \cdots t_n = s$. Let $\Delta_1, \ldots, \Delta_q$ be the domains of transitivity of the cyclic group $\langle s \rangle$ where $\# \Delta_i = e_i \geq 1$. Following [Mo] pp.369,370 if $s = s_1 \cdots s_q$ is the corresponding product of independent cycles we may write $s_i = (1_i 2_i \ldots (e_i)_i)$ if $e_i > 1$ and $s_i = (1_i) = 1$ if $e_i = 1$. Such a representation is uniquely determined by s if we assume that that 1_i is the minimal number among Δ_i for each i and if we order Δ_i in such a way that $1_1 < 1_2 < \cdots < 1_q$. Let $e_i > 1$ and let $\mathcal{Z}_i$ be the sequence $((1_i 2_i), (1_i 3_i), \ldots, (1_i (e_i)_i))$. If $e_i = 1$ one lets $\mathcal{Z}_i = \emptyset$. Let $\mathcal{Z}$ be the concatenation of sequences $\mathcal{Z} = \mathcal{Z}_1 \mathcal{Z}_2 \ldots \mathcal{Z}_q$. The sequence $\mathcal{Z}$ consists of $N = \sum_{i=1}^{q} (e_i - 1)$ transpositions whose product equals s. The following theorem is proved in [Mo] pp.369,370. It can also be deduced from the earlier paper of Kluitmann [Kl].

Theorem 3.5. *Given a sequence of transpositions* $\underline{t} = (t_1, \ldots, t_n)$ *satisfying* $t_1 \cdots t_n = s$ *and such that* $\langle t_1, \ldots, t_n \rangle$ *is a transitive group (therefore* $= S_d$*), there is a braid-equivalent sequence* $(t'_1, \ldots, t'_n)$ *which has the following form*

$$(t'_1, \ldots, t'_n) \; = \; (\mathcal{Z}, t'_{N+1}, \ldots, t'_n)$$

where $n - N \equiv 0 (mod\, 2)$ *and*

1. *if $q = 1$, then $t'_i = (1_1 2_1)$ for $\forall i \geq N+1$*
2. *if $q > 1$ then*

$$(t'_{N+1}, \ldots, t'_n) \;=\; ((1_1 1_2), (1_1 1_2), (1_1 1_3), (1_1 1_3), \ldots, (1_1 1_q), (1_1 1_q))$$

where each $(1_1 1_i)$ appears two times if $2 \leq i \leq q-1$ and $(1_1 1_q)$ appears an even number of times .

Corollary 3.6. *Let $\underline{t} = (t_1, \ldots, t_n)$ be a sequence of transpositions in S_d which generate a transitive group. Let $s = t_1 \cdots t_n$ and let $\Delta_1, \ldots, \Delta_q$ be the domains of transitivity of the cyclic group $\langle s \rangle$. Suppose that $q \geq 2$. Let $a \in \Delta_i$, $b \in \Delta_j$ where $i \neq j$. Then $\underline{t}$ is braid-equivalent to a sequence $(\ldots, (ab), (ab))$.*

Proof. Let us consider the group S_q which permutes the indices of Δ_i, $i = 1, \ldots, q$. Let $\tau_1 = (ij), \tau_2, \ldots, \tau_{q-1}$ be transpositions in S_q which generate it. We let $a_i = a$, $a_j = b$ and choose arbitrarily $a_k \in \Delta_k$ for each $k \neq i, j$. If $\tau_\ell = (\alpha, \beta)$ we consider the pair of transpositions $(a_\alpha a_\beta), (a_\alpha a_\beta)$. We then consider a sequence of n transpositions in S_d which is a concatenation of the sequence $\mathscr{Z}$, of the pairs associated with $\tau_1, \tau_2, \ldots, \tau_{q-1}$ and of the sequence $(1\,2), \ldots, (1\,2)$ with (12) repeated an even number of times. We obtain a sequence $\mathscr{Z}, (ab), (ab), \ldots$ with product s which generates a transitive group. According to Theorem 3.5 this sequence is braid-equivalent to $\underline{t}$. Moving the pair $(ab), (ab)$ to the end we obtain the required sequence. ☐

3.7. Given a permutation $s \in S_d$ whose domains of transitivity are $\Delta_1, \ldots, \Delta_q$ with $\#\,\Delta_i = e_i$ we let $|s| = \sum_{i=1}^{q}(e_i - 1)$. Let us consider a sequence of transpositions $\underline{t} = (t_1, \ldots, t_n)$, $t_i \in S_d$. Let $\underline{\Sigma} = \{\Sigma_1, \ldots, \Sigma_m\}$ be the domains of transitivity of the group $\langle t_1, \ldots, t_n \rangle$. We let $|\underline{\Sigma}| = \sum_{j=1}^{m}(\#\,\Sigma_j - 1)$.

Lemma 3.8. *Let $\underline{t} = (t_1, \ldots, t_n)$ be a sequence of transpositions and let $t_1 \cdots t_n = s$. Then $n + |s| \equiv 0 (mod\, 2)$ and $n + |s| \geq 2|\underline{\Sigma}|$. If $n + |s| > 2|\underline{\Sigma}|$ then there is a sequence $\underline{t}'$ braid-equivalent to $\underline{t}$ such that $t'_{n-1} = t'_n$ and $t'_1, \ldots, t'_{n-2}$ generate the same group as $t_1, \ldots, t_n$.*

Proof. It is clear that every Δ_i belongs to some Σ_j. Furthermore one may replace $\underline{t}$ by a braid-equivalent sequence which is a concatenation $T_1 T_2 \ldots T_m$ of sequences T_j whose elements belong to $S(\Sigma_j)$ (cf. the proof of Lemma 3.3; if $\#\,\Sigma_j = 1$ one lets $T_j = \emptyset$). It thus suffices to prove the lemma in the case when $\langle t_1, \ldots, t_n \rangle$ is transitive, which we further assume, so $\underline{\Sigma} = \{\Sigma_1\}$, $|\underline{\Sigma}| = d - 1$. Let us consider the braid-equivalent sequence $\underline{t}'$ of Theorem 3.5. The subsequence $\mathscr{Z}$ has $N = |s|$ elements. The remaining $n - |s|$ elements appear in pairs and their number is at least $\geq 2(q-1)$ since $\langle t'_1, \ldots, t'_n \rangle$ is a transitive group. If $n - |s| > 2(q-1)$ the pair $(1_1 1_q), (1_1 1_q)$ is repeated at least twice, so canceling it from the sequence does not change the group generated by the transpositions t'_i. One has $\Sigma_1 = \cup_{i=1}^{q} \Delta_i$, so $d = |s| + q$. Adding $2|s|$ to both sides of the inequality $n - |s| \geq 2(q-1)$ we obtain the equivalent inequality $n + |s| \geq 2d - 2 = 2|\Sigma_1|$. ☐

Definition 7. A transitive subgroup $G \subset S_d$ is called *imprimitive* if there is a decomposition $\{1, \dots, d\} = \bigsqcup_{i=1}^{k} \Sigma_i$ such that for $\forall g \in G$ and $\forall i$ one has $\Sigma_i{}^g = \Sigma_j$ for some j and furthermore $\# \Sigma_i = e$ for $\forall i$ where $1 < e < d$. A transitive subgroup $G \subseteq S_d$ which is not imprimitive is called *primitive.*

If d is a prime number clearly every transitive subgroup of S_d is primitive.

Lemma 3.9. *Suppose that $G \subseteq S_d$ is a primitive group which contains a transposition. Then $G = S_d$. In particular if d is a prime number every transitive subgroup of S_d which contains a transposition is equal to S_d.*

Proof. Let us prove that if G contains a nontrivial symmetric subgroup $S(\Sigma)$ with $\# \Sigma < d$ then G contains a $S(\Sigma')$ with $\# \Sigma' = \# \Sigma + 1$. Indeed, since G is primitive there are $g, h \in G$ such that $0 < \#(\Sigma^g \cap \Sigma^h) < \# \Sigma$. Hence if $f = hg^{-1}$ one has $0 < \#(\Sigma \cap \Sigma^f) < \# \Sigma$. Let $x^f \in \Sigma \cap \Sigma^f$, $y^f \in \Sigma^f - \Sigma$. Then $x, y \in \Sigma$, $x^f \in \Sigma$, $y^f \notin \Sigma$. So $\Sigma' = \Sigma \cup \{x, y\} \supsetneqq \Sigma$ and $S(\Sigma') = \langle S(\Sigma), f^{-1}(xy)f \rangle$ is contained in G. We conclude the proof of the lemma by induction on $\# \Sigma$ starting from $\Sigma = \{a, b\}$ with $(ab) \in G$. □

To each connected component of $\mathcal{H}_{d,n}(Y, b_0)$ is associated a fixed monodromy group $G \subseteq S_d$ and similarly to each connected component of $\mathcal{H}_{d,n}(Y)$ is associated a conjugacy class of subgroups.

Definition 8. Let $d \geq 2$, $n > 0$. We denote by $\mathcal{H}^0_{d,n}(Y, b_0)$ and $\mathcal{H}^0_{d,n}(Y))$ the union of connected components of $\mathcal{H}^{(n)}_d(Y, b_0)$ and $\mathcal{H}^{(n)}_d(Y)$ respectively which correspond to coverings with primitive monodromy groups (therefore equal to S_d by Lemma 3.9).

The following theorem is due in the case $n \geq 2d$ to Graber, Harris and Starr [GHS].

Theorem 3.10. *Let Y be a smooth, projective curve. Let $b_0 \in Y$. If $n \geq 2d - 2$ the Hurwitz spaces $\mathcal{H}^0_{d,n}(Y, b_0)$ and $\mathcal{H}^0_{d,n}(Y)$ are irreducible.*

Proof. The case $Y \cong \mathbb{P}^1$ is classical and due to Clebsch and Hurwitz [Cl], [Hu]. Modern proofs may be found in [Mo] p.368 and [Vo] p.197. Suppose that $g(Y) \geq 1$. Let us first consider $\mathcal{H}^0_{d,n}(Y, b_0)$. Since $\mathcal{H}^0_{d,n}(Y, b_0)$ is smooth in order to prove its irreducibility it suffices to prove it is connected. By Theorem 1.10 it suffices to show that every Hurwitz system $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ with monodromy group S_d is braid-equivalent to

$$((12), (12), (13), (13), \dots, (1d), (1d); 1, 1, \dots, 1, 1) \tag{11}$$

where each $(1i)$ appears twice if $2 \leq i \leq d - 1$ and $(1d)$ appears $n - 2d + 4$ times.

Step 1. We claim that every Hurwitz system $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ is braid-equivalent to a Hurwitz system $(\tilde{t}_1, \dots, \tilde{t}_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ such that $\langle \tilde{t}_1, \dots, \tilde{t}_n \rangle$ is a transitive group. Indeed, let $s = [\lambda_1, \mu_1] \cdots [\lambda_g, \mu_g]$ and let $\underline{\Sigma} = \{\Sigma_1, \dots, \Sigma_m\}$ be the domains of transitivity of the group $\langle t_1, \dots, t_n \rangle$. One has $|\underline{\Sigma}| = d - m$, so if $\langle t_1, \dots, t_n \rangle$ is not transitive then $|\underline{\Sigma}| < d - 1$. In this case the inequality $n + |s| > 2|\underline{\Sigma}|$ is satisfied, so according to

Lemma 3.8 one may replace $(t_1,\dots,t_n)$ by a braid-equivalent sequence $(t'_1,\dots,t'_n)$ such that $t'_{n-1}=t'_n=(ab)$ and $\langle t'_1,\dots,t'_{n-2}\rangle=\langle t_1,\dots,t_n\rangle$, so $H=\langle t'_1,\dots,t'_{n-2};\lambda_1,\mu_1,\dots,\lambda_g,\mu_g\rangle=S_d$. Suppose that $a,b\in\Sigma_i$. Let $h\in H$ be such that $a^h\in\Sigma_1,\ b^h\notin\Sigma_1$. Then according to Main Lemma 2.1 the Hurwitz system $(t'_1,\dots,t'_{n-2},t'_{n-1},t'_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$ is braid-equivalent to $(t'_1,\dots,t'_{n-2},(t'_{n-1})^h,(t'_n)^h;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$. If $\underline{\Sigma}'=\{\Sigma'_1,\dots,\Sigma'_r\}$ are the domains of transitivity of $\langle t'_1,\dots,t'_{n-2},(t'_{n-1})^h,(t'_n)^h\rangle$ then clearly $|\underline{\Sigma}'|>|\underline{\Sigma}|$. Repeating this argument after a finite number of steps one obtains a braid-equivalent Hurwitz system $(\tilde t_1,\dots,\tilde t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$ with transitive $\langle\tilde t_1,\dots,\tilde t_n\rangle$.

Step 2. Let $(t_1,\dots,t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$ be a Hurwitz system. Suppose that $\sum_{\ell=1}^g(|\lambda_\ell|+|\mu_\ell|)>0$. We claim there is a braid-equivalent sequence $(t'_1,\dots,t'_n;\lambda'_1,\mu'_1,\dots,\lambda'_g,\mu'_g)$ such that $\sum_{\ell=1}^g(|\lambda'_\ell|+|\mu'_\ell|)<\sum_{\ell=1}^g(|\lambda_\ell|+|\mu_\ell|)$. Let us first suppose that $\lambda_1\neq1$. Decomposing λ_1 into a product of nontrivial independent cycles and choosing one of them $(ab\dots c)$ we have $|(ab)\lambda_1|<|\lambda_1|$. According to Step 1 we may replace $(t_1,\dots,t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$ by a braid-equivalent system $(\tilde t_1,\dots,\tilde t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$ such that $\tilde t_1,\dots,\tilde t_n$ generate a transitive group. Then according to Mochizuki's Lemma 3.3 we may replace $(\tilde t_1,\dots,\tilde t_n)$ by a braid-equivalent sequence $((ab),t'_2,\dots,t'_n)$. Applying the braid move τ''_{11} (cf. Eq.(10)) we transform λ_1 into $(ab)\lambda_1$. Suppose that $\lambda_1=1,\ \mu_1\neq1$. We reason in the same way and use the braid move ρ'_{11} which transforms $((ab),\dots;1,\mu_1,\dots)$ into $(t'_1,\dots;1,(ab)\mu_1,\dots)$ according to Corollary 1.9 (i). Similarly if $\lambda_k\neq1$ and $\lambda_\ell=\mu_\ell=1$ for $\forall\ell\le k-1$ we have $u_{k-1}=1$ and applying the braid move τ''_{1k} we transform $((ab),\dots;\dots,\lambda_k,\mu_k,\dots)$ into $(t'_1,\dots;\dots,(ab)\lambda_k,\mu_k,\dots)$ thus decreasing $|\lambda_k|$. If $\mu_k\neq1$, $\lambda_\ell=\mu_\ell=1$ for $\forall\ell\le k-1$ and $\lambda_k=1$ we apply ρ'_{1k} to transform $((ab),\dots;\dots,1,\mu_k,\dots)$ into $(t'_1,\dots;\dots,1,(ab)\mu_k,\dots)$ thus decreasing $|\mu_k|$.

Step 3. Starting with $(t_1,\dots,t_n;\lambda_1,\mu_1,\dots,\lambda_g,\mu_g)$ and applying several times Step 2 one obtains a braid-equivalent Hurwitz system $(\tilde t_1,\dots,\tilde t_n;1,1,\dots,1,1)$. Here $\tilde t_1\cdots\tilde t_n=1$, so applying the argument in the case $g(Y)=0$ (see e.g. [Mo] p.368, or [Vo] p.197) one obtains the initial Hurwitz system is braid-equivalent to Eq.(11).

The connectedness of both $\mathcal{H}^0_{d,n}(Y,b_0)$ and $\mathcal{H}^0_{d,n}(Y)$ follows now from Theorem 1.10. □

Theorem 3.10 may be generalized in a straightforward manner to coverings $\pi:X\to Y$ which have simple ramifications in all except possibly one discriminant point. The conjugacy classes of S_d correspond bijectively to partitions of d, namely $\underline{e}=\{e_1,e_2,\dots,e_q\}$ where $e_1\ge e_2\ge\ldots\ge e_q\ge1$ and $e_1+\cdots+e_q=d$. To each partition one associates the orbit of the permutation

$$\epsilon=(1\,2\dots e_1)(e_1+1\dots e_1+e_2)\dots(\sum_{i=1}^{q-1}e_i\ +1\dots d)\tag{12}$$

Given a partition $\underline{e}$ let us denote by $\mathcal{H}_{d,n,\underline{e}}(Y,b_0)$ and $\mathcal{H}_{d,n,\underline{e}}(Y))$ the Hurwitz spaces which parametrize equivalence classes $[X\to Y,\,\phi]$ and $[X\to Y]$ respectively, where X is connected and $\pi:X\to Y$ is a covering of degree d with n discriminant points of simple ramification and one additional discriminant point whose local monodromy belongs to the conjugacy class of Eq.(12). Let $\mathcal{H}^0_{d,n,\underline{e}}(Y,b_0)$ and $\mathcal{H}^0_{d,n,\underline{e}}(Y)$ parametrize such coverings with primitive monodromy group.

Theorem 3.11. *Let Y be a smooth, projective curve. Let $b_0 \in Y$. Suppose that $n \geq 2d-2$. Let $\underline{e}$ be an arbitrary partition of d. Then the Hurwitz spaces $\mathcal{H}^0_{d,n,\underline{e}}(Y,b_0)$ and $\mathcal{H}^0_{d,n,\underline{e}}(Y)$ are irreducible.*

Proof. If $\underline{e}$ is the trivial partition (that is $e_i = 1$ for $\forall i$) this is the content of Theorem 3.10. Suppose that $e_1 \geq 2$. Let us denote the permutation Eq.(12) by $\epsilon = (1_1 2_1 \ldots (e_1)_1)(1_2 \ldots (e_2)_2)\ldots$. The theorem will be proved if we can show that every Hurwitz system $(t_1,\ldots,t_n,t_{n+1};\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ of the considered type may be reduced after a finite number of braid moves of the types $\sigma'_j, \sigma''_j, \rho'_{ik}, \rho''_{ik}, \tau'_{ik}, \tau''_{ik}$ to the normal form

$$(\mathcal{Z}, t'_{N+1},\ldots,t'_n,\epsilon^{-1};1,1,\ldots,1,1)$$

where the n-tuple of transpositions $(\mathcal{Z}, t'_{N+1},\ldots,t'_n)$ is the one defined in 3.4 and Theorem 3.5. First using braid moves of type σ'_j we may replace the original Hurwitz system by one for which t_{n+1} belongs to the orbit of ϵ. We then let $s = [\lambda_1,\mu_1]\cdots[\lambda_g,\mu_g]t^{-1}_{n+1}$ and repeating the arguments of the proof of Theorem 3.10 we obtain a braid-equivalent Hurwitz system $(\tilde{t}_1,\ldots,\tilde{t}_n,\tilde{t}_{n+1};1,1,\ldots,1,1)$ with $\tilde{t}_{n+1} = t_{n+1}$. We are further allowed to apply only braid moves of types σ'_j and σ''_j. The next step is to replace the obtained Hurwitz system by one in which at the $(n+1)$-th place stays ϵ^{-1}. In fact the permutation ϵ^{-1} has the same cyclic type as $\tilde{t}_{n+1}$, so $\epsilon^{-1} = a^{-1}\tilde{t}_{n+1}a$. Let $a = \tau_1\cdots\tau_r$ where τ_i are transpositions. By the hypothesis and Lemma 3.9 we have $\langle \tilde{t}_1,\ldots,\tilde{t}_n\rangle = S_d$. So using Mochizuki's Lemma 3.3 we may replace $(\tilde{t}_1,\ldots,\tilde{t}_n)$ by a braid-equivalent n-tuple $(\ldots,\tau_1)$. We then have $(\tau_1,\tilde{t}_{n+1}) \sim (\tau_1\tilde{t}_{n+1}\tau_1,\tau_1) \sim (\tau'_1,\tau_1\tilde{t}_{n+1}\tau_1)$, so $(\tilde{t}_1,\ldots,\tilde{t}_n,\tilde{t}_{n+1}) \sim (\ldots,\tau_1\tilde{t}_{n+1}\tau_1)$. Repeating this argument with $\tau_2,\tau_3,\ldots,\tau_r$ we obtain a braid-equivalent $(n+1)$-tuple whose $(n+1)$-th element is ϵ^{-1} and the product of the first n equals ϵ. Using a finite number of braid moves among the first n transpositions we obtain the required normal form according to Theorem 3.5. ☐

So far we worked with coverings with primitive monodromy groups. We now wish to consider the imprimitive case.

Lemma 3.12. *Let $G \subset S_d$ be a transitive imprimitive subgroup which contains a transposition. Then there is a unique decomposition $\{1,\ldots,d\} = \Sigma_1 \sqcup \ldots \sqcup \Sigma_k$ as in Definition 7 so that $G_i = G \cap S(\Sigma_i)$ is a primitive subgroup of $S(\Sigma_i)$ for $\forall i$. Furthermore the following properties hold.*

i. All transpositions contained in G form one conjugacy G-orbit T.

ii. $G_1 \cdot G_2 \cdots G_k \;=\; G_1 \times G_2 \times \cdots \times G_k = \langle T\rangle$.

iii. $G_i = S(\Sigma_i)$ for $\forall i$.

Proof. Let $(ab) \in G$ and let $T = (ab)^G$. Let $H = \langle T \rangle$ and let $\Sigma_1, \dots, \Sigma_k$ be the domains of transitivity of H. Since H is a normal subgroup every $g \in G$ permutes the orbits $\{\Sigma_1, \dots, \Sigma_k\}$. Furthermore $H = S(\Sigma_1) \times \cdots \times S(\Sigma_k)$, so $G_i = G \cap S(\Sigma_i) = S(\Sigma_i)$. All transpositions which belong to H form one G-orbit since this property holds for $S(\Sigma_i)$. Let $(\alpha\,\beta)$ be an arbitrary transposition of G. We claim it is impossible that $\alpha \in \Sigma_i,\ \beta \in \Sigma_j$ for $i \neq j$. Indeed, if this were the case letting $g = (\alpha\,\beta)$ we would have ${\Sigma_i}^g \cap \Sigma_j \neq \emptyset$, so ${\Sigma_i}^g = \Sigma_j$. If $\alpha' \in \Sigma_i, \alpha' \neq \alpha$, then $\alpha'^g = \alpha' \in \Sigma_i$ and on the other hand $\alpha'^g \in \Sigma_j$, so $\Sigma_i \cap \Sigma_j \neq \emptyset$ which is absurd. We obtain $\{\alpha\,\beta\} \subseteq \Sigma_i$ for some i, so $(\alpha\,\beta) \in T = (ab)^G$. This proves (i). Suppose now that $\{1, \dots, d\} = \Sigma'_1 \sqcup \ldots \sqcup \Sigma'_\ell$ is an arbitrary decomposition as required in Definition 7. Let $(\alpha\,\beta)$ be a transposition in G. Then the argument above shows that $\{\alpha, \beta\} \subseteq \Sigma'_i$ for some i. This shows that $T = (\alpha\,\beta)^G \subseteq G' = G'_1 \times \cdots \times G'_\ell$ where $G'_j = G \cap S(\Sigma'_j)$. Thus each orbit Σ_i of $H = \langle T \rangle$ is contained in some orbit Σ'_j. Assuming G'_j is primitive subgroup of $S(\Sigma'_j)$ we conclude that $\Sigma_i = \Sigma'_j$. This proves the uniqueness statement. The other properties were already proved. ☐

Corollary 3.13. *Let $\pi : X \to Y$ be a covering of smooth, irreducible, projective curves of degree d. Suppose that at least one of the discriminant points is simple and suppose that the monodromy group of the covering is $\neq S_d$. Then there exists a unique decomposition $X \xrightarrow{\pi_1} \tilde{Y} \xrightarrow{\pi_2} Y$, $\pi = \pi_2 \circ \pi_1$, where $d_i = \deg \pi_i > 1$ for $i = 1, 2$ and $\pi_1 : X \to \tilde{Y}$ has primitive monodromy group. If $\pi : X \to Y$ has n simple discriminant points so does $\pi_1 : X \to \tilde{Y}$ and π_2 induces a bijective correspondence between these two sets.*

Proof. One chooses an unramified point $b_0 \in Y$, and a bijection $\phi : \pi^{-1}(b_0) \longrightarrow \{1, \dots, d\}$. One applies Lemma 3.12 to the monodromy group of $X \to Y$ and obtains a decomposition $X \to \tilde{Y} \to Y$. Replacing ϕ by another bijection results in replacing the monodromy group G by a conjugate one $G' = s^{-1}Gs$. The conjugation by $s \in S_d$ transforms the G-orbit of transpositions contained in G to the G'-orbit of transpositions contained in G'. Therefore s transforms the corresponding decompositions of $\{1, \dots, d\}$: $\Sigma'_i = {\Sigma_i}^s$. This proves the uniqueness of the decomposition $X \to \tilde{Y} \to Y$ with the required properties. The last statement of the corollary is obvious. ☐

Theorem 3.14. *Let Y be a smooth, projective curve of genus ≥ 1. Let $d \geq 2$, $n > 0$, $n \equiv 0 (mod\ 2)$. If d is prime then $\mathcal{H}_{d,n}(Y) = \mathcal{H}^0_{d,n}(Y)$. If d is not prime let d' be its maximal divisor $\neq d$. Suppose that $n \geq 2d' - 2$. Then the connected components of $\mathcal{H}_{d,n}(Y)$ which correspond to simple coverings with monodromy groups $\neq S_d$ are in one-to-one correspondence with the equivalence classes of unramified coverings $[\tilde{Y} \to Y]$ of degrees $d_2 | d$ where $d_2 \neq 1, d$.*

Proof. If d is prime every transitive subgroup of S_d is primitive, so $\mathcal{H}_{d,n}(Y) = \mathcal{H}^0_{d,n}(Y)$. Suppose that d is not prime. Let $\pi : X \to Y$ be a simple covering of degree d, simply ramified in n points, with monodromy group $\neq S_d$. Then the decomposition $X \xrightarrow{\pi_1} \tilde{Y} \xrightarrow{\pi_2} Y$ of Corollary 3.13 is with étale π_2. Furthermore by the uniqueness of this decomposition if $f : X \to X'$ defines an equivalence of coverings of Y then f is induced by an equivalence $g : \tilde{Y} \to \tilde{Y}'$. Given an étale covering $p_2 : \tilde{Y} \to Y$ of degree $d_2 \neq 1, d$ we consider the Hurwitz space $\mathcal{H}^0_{d_1,n}(\tilde{Y})$ where $d_1 = d/d_2$. It is connected by Theorem 3.10. We then consider the Zariski open subset consisting of $[X \to \tilde{Y}]$ with discriminant points belonging to different fibers of $\pi_2 : \tilde{Y} \to Y$. Every such $X \to \tilde{Y}$ yields a simple covering of Y. In this way one obtains a connected component of $\mathcal{H}_{d,n}(Y)$ whose conjugacy class of monodromy groups is imprimitive. Vice versa by Corollary 3.13 every connected component of $\mathcal{H}_{d,n}(Y)$ whose associated conjugacy class of monodromy groups is imprimitive is obtained in this way. $\square$

Example. Let $d = 4$ and let Y be a curve of genus $g \geq 1$. The connected unramified coverings of Y of degree 2 are classified up to equivalence by the points of order 2 in the Jacobian $J(Y)$. Applying Theorem 3.14 we obtain if $n \geq 2$ there are $2^{2g} - 1$ different connected components of $\mathcal{H}_{4,n}(Y)$ which parametrize coverings with imprimitive monodromy group. The monodromy group in this case is isomorphic to the dihedral group D_4

4 The case $n = 2d - 4, \quad g \geq 1$

Unless otherwise specified in this section we shall work with Hurwitz systems $(t_1, \ldots, t_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$ where $n \geq 2,\ g \geq 1$ and $t_1, \ldots, t_n$ are transpositions in S_d. Let $T = (t_1, \ldots, t_n)$ be a sequence of transpositions in S_d and let $s = t_1 \cdots t_n$. Suppose that the subgroup $\langle T \rangle = \langle t_1, \ldots, t_n \rangle$ has only one nontrivial domain of transitivity Σ and let $e = \#\,\Sigma$. We have the inequality $n \geq e - 1,\ n + |s| \geq 2(e-1)$ (cf. Lemma 3.8) so if $e = \#\,\Sigma$ is fixed the minimum for n is reached for $n = e - 1$, and then $|s| = e - 1$. This happens if and only if $s = t_1 \cdots t_n$ is a cycle of order $n + 1$.

Definition 9. We call a sequence of transpositions $T = (t_1, \ldots, t_n)$ *minimal* if $s = t_1 \cdots t_n$ is a cycle of order $n + 1$.

Lemma 4.1. *Let $T = (t_1, \ldots, t_n)$ be a sequence of transpositions of S_d such that $\langle T \rangle = \langle t_1, \ldots, t_n \rangle$ has a unique nontrivial domain of transitivity Σ. Suppose that T is not minimal. Then for any $a \in \Sigma$ there exists a braid-equivalent sequence $T' = (t'_1, \ldots, t'_n)$ such that $t'_{n-1} = t'_n = (ab)$ for some $b \in \Sigma,\ b \neq a$.*

Proof. Without loss of generality we may assume that $\Sigma = \{1, \ldots, d\}$. Let $s = t_1 \cdots t_n$ and let $\Delta_1, \ldots, \Delta_q$ be the domains of transitivity of s. Replacing $(t_1, \ldots, t_n)$ by a braid-equivalent sequence we may assume it has the normal form of Theorem 3.5. The non-minimality hypothesis $n > |s|$ implies that T contains $(n - |s|)/2$ pairs of the form $(1_1 2_1)$ if $q = 1$ or $(1_1 1_i)$ with $2 \leq i \leq q$ if $q \geq 2$. Consider the cyclic group $\langle s \rangle$. If $u \in \langle s \rangle$ then ${t_1}^u \cdots {t_n}^u = s$, so the sequence $({t_1}^u, \ldots, {t_n}^u)$ is braid-equivalent to $(t_1, \ldots, t_n)$ (cf. [Kl], [Mo]). If $q = 1$ we may find $u \in \langle s \rangle$ such that $(1_1)^u = a$. Here we let $b = (2_1)^u$. If $q \geq 2$ and $a \in \Delta_1$ we find u such that $(1_1)^u = a$ and we let $b = (1_2)^u$. If $a \in \Delta_i$ with $i \geq 2$ we find $u \in \langle s \rangle$ such that $(1_i)^u = a$ and we let $b = (1_1)^u$. In each case the sequence $({t_1}^u, \ldots, {t_n}^u)$ contains the adjacent pair $(ab), (ab)$. Moving it to the end we obtain the sequence required in the lemma. $\square$

4.2. Given a Hurwitz system $(t_1,\ldots,t_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ where $t_i,\lambda_k,\mu_k\in S_d$ let $\underline{\Sigma}=\{\Sigma_1,\Sigma_2,\ldots,\Sigma_m\}$ be the domains of transitivity of $G_1=\langle t_1,\ldots,t_n\rangle$ ordered in such a way that $\#\,\Sigma_1\geq\#\,\Sigma_2\geq\cdots\geq\#\,\Sigma_m$. Let

$$\ell(\underline{\Sigma}) \;=\; (\#\,\Sigma_1,\#\,\Sigma_2,\ldots,\#\,\Sigma_m,0,0,\ldots) \tag{13}$$

be the associated partition of d. Given Σ_i let T_i be the subsequence of $T=(t_1,\ldots,t_n)$ composed of those transpositions which move points of Σ_i. Let $n_i=\#\,T_i$. We call Σ_i minimal or non-minimal if the sequence T_i is minimal or non-minimal respectively. The trivial case $\#\,\Sigma_i=1,\ T_i=\emptyset$ is assumed minimal.

Lemma 4.3. *Let $(t_1,\ldots,t_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ be a Hurwitz system with monodromy group S_d. Let $s=[\lambda_1,\mu_1]\cdots[\lambda_g,\mu_g]$ and let $G_1=\langle t_1,\ldots,t_n\rangle\subseteq S_d$. Let $\underline{\Sigma}=\{\Sigma_1,\ldots,\Sigma_m\}$ be the domains of transitivity of G_1 ordered as in 4.2. Suppose that $\ell(\underline{\Sigma})$ is maximal in lexicographic order among the partitions associated with all Hurwitz systems of the type $(t'_1,\ldots,t'_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ braid-equivalent to the given one. Then $\underline{\Sigma}=\{\Sigma_1\}$ (equivalently $\langle t_1,\ldots,t_n\rangle$ is transitive) if and only if $n+|s|\geq 2d-2$. If $n+|s|<2d-2$ then $n+|s|=2|\underline{\Sigma}|=2(d-m)$. Furthermore $n_1+|s|_{\Sigma_1}|=2(\#\,\Sigma_1-1)$ and for each $i\geq 2$ the G_1-orbit Σ_i is minimal, equivalently $n_i=|s|_{\Sigma_i}|=\#\,\Sigma_i-1$. In particular if $s=1$ then $\langle t_1,\ldots,t_n\rangle$ has a single nontrivial domain of transitivity.*

Proof. We proved in Lemma 3.8 that if $\langle t_1,\ldots,t_n\rangle$ is transitive then $n+|s|\geq 2d-2$. That the maximality of $\ell(\underline{\Sigma})$ and $n+|s|\geq 2d-2$ imply that $\underline{\Sigma}=\{\Sigma_1\}$ is a fact evident from the proof of Theorem 3.10 (cf. Step 1). We shall further assume that $n+|s|<2d-2$, so G_1 is not transitive. Suppose, by way of contradiction, that there is a non-minimal T_k with $k\geq 2$. Let $\Sigma'=\Sigma_{k+1}\cup\cdots\cup\Sigma_m$, so $\{1,\ldots,d\}=\Sigma_1\cup\cdots\cup\Sigma_k\cup\Sigma'$. Let $H_1=\langle T_k\rangle$, $H_2=\langle T_1,\ldots,\widehat{T_k},\ldots,T_m,\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g\rangle$, so $\langle H_1,H_2\rangle=G=S_d$. For every $b\in\Sigma_k$ one has $b^G=\{1,\ldots,d\}$. Let $a\in\Sigma_k$ and $h\in G=\langle H_1,H_2\rangle$ be chosen in such a way that $a^h\in\Sigma_1\cup\cdots\cup\Sigma_{k-1}$ and furthermore the length of the factorization $h=h_1h_2\cdots h_r$ with $h_i\in H_1$ or H_2 is minimal possible among all $b\in\Sigma_k$ and all $g\in G$ satisfying $b^g\in\Sigma_1\cup\cdots\cup\Sigma_{k-1}$. We claim $r=1$ and $h_1\in H_2$. In fact by minimality we have $h_1\in H_2$, $h_r\in H_2$ and for each $i=1,\ldots,r-1$ the adjacent h_i,h_{i+1} do not belong to the same subgroup H_1 or H_2. Suppose that $r\geq 2$. Then by minimality $a^{h_1\cdots h_{r-1}}\in\Sigma_k\cup\Sigma'$. If $a^{h_1\cdots h_{r-1}}\in\Sigma_k$ we might replace a by $a'=a^{h_1\cdots h_{r-1}}$, so $a^{h_1\cdots h_{r-1}}\in\Sigma'$. The group H_1 acts trivially on Σ', therefore since $h_{r-1}\in H_1$ we have $a^{h_1\cdots h_{r-1}}=a^{h_1\cdots h_{r-2}}$. Hence $a^h=a^{h_1\cdots(h_{r-2}h_r)}=a^{h'}$ where h' has a shorter factorization then h. This is an absurd, so $r=1$. We have thus proved there exists an $a\in\Sigma_k$ and $h\in H_2$ such that $a^h\in\Sigma_i$ for some $i<k$. According to Lemma 4.1 we may replace T_k by a braid-equivalent sequence which contains the adjacent pair $(ab),(ab)$ for some $b\in\Sigma_k$. By Main Lemma 2.1 replacing $(ab),(ab)$ by $(ab)^h,(ab)^h$ one obtains a braid-equivalent Hurwitz system. The following cases may occur.

Case 1. *$b^h\in\Sigma_j$ for some $j\neq i,\ 1\leq j\leq m$.* If $i<j$ then the new Hurwitz system has $\Sigma_i\cup\Sigma_j$ as a domain of transitivity, so the corresponding partition Eq.(13) is greater than $\ell(\underline{\Sigma})$ in lexicographical order. If $j<i$ we interchange a and b and make the same conclusion. In both cases we obtain a contradiction with the maximality of $\ell(\underline{\Sigma})$.

Case 2. $b^h \in \Sigma_i$. Let $(ab)^h = (xy)$. Moving the pair $(xy),(xy)$ to the right of T_i we obtain a braid-equivalent sequence $\tilde{T}_i = T_i,(xy),(xy)$ with the same domain of transitivity as T_i. Canceling the pair from the obtained Hurwitz system does not change the monodromy group G. Let us choose $g \in G = S_d$ such that $x^g \in \Sigma_i$, $y^g \notin \Sigma_i$. Applying Main Lemma 2.1 we obtain a new Hurwitz system $(\tilde{t}_1,\ldots,\tilde{t}_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ whose associated partition $\ell(\underline{\Sigma})$ has the property that its first i terms are greater in lexicographical order then the first i terms of $\ell(\underline{\Sigma})$. This contradicts the maximality of the partition $\ell(\underline{\Sigma})$.

The claim that T_i is minimal for every $i \geq 2$ is proved. According to Lemma 3.8 we have $n_1 + |s|_{\Sigma_1}| \geq 2(\#\Sigma_1 - 1)$ and if this inequality is strict we can apply the argument of Step 1 of Theorem 3.10 in order to increase $\ell(\underline{\Sigma})$ which is impossible . Thus $n_1 + |s|_{\Sigma_1}| = 2(\#\Sigma_1 - 1)$. Finally if $s = 1$ then $s|_{\Sigma_i} = 1$, so minimality of Σ_i means $\#\Sigma_i = 1$. Therefore the maximality of $\ell(\underline{\Sigma})$ implies that only Σ_1 may be nontrivial domain of transitivity. □

4.4. Given a Hurwitz system $(t_1,\ldots,t_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ we let $\underline{\lambda} = (\lambda_1,\ldots,\lambda_g)$, $\underline{\mu} = (\mu_1,\ldots,\mu_g)$, $|\underline{\lambda}| = (|\lambda_1|,\ldots,|\lambda_g|)$, $|\underline{\mu}| = (|\mu_1|,\ldots,|\mu_g|)$ (cf. 3.7). We consider the class $\mathcal{C}$ of Hurwitz systems braid-equivalent to the given one. When dealing with the problem of finding a system in $\mathcal{C}$ of a simplest form it makes sense to assume (without loss of generality) that $(t_1,\ldots,t_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ satisfies the following property:

(*) $(|\underline{\lambda}|,|\underline{\mu}|)$ *is minimal in lexicographic order among the Hurwitz systems in* $\mathcal{C}$ *and furthermore* $\ell(\underline{\Sigma})$ *(cf. 4.2) is maximal in lexicographic order among the Hurwitz systems in* $\mathcal{C}$ *of the type* $(t'_1,\ldots,t'_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$

Theorem 4.5. *Let* $d \geq 3$, $n = 2d-4$. *Let* Y *be a smooth, projective curve of genus* $g \geq 1$. *Then the Hurwitz space* $\mathcal{H}^0_{d,n}(Y)$ *parametrizing simple coverings with monodromy groups equal to* S_d *is irreducible.*

Proof. By Theorem 1.10 it suffices to show that the equivalence class (modulo inner automorphisms) of every Hurwitz system $(t_1,\ldots,t_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ with monodromy group S_d may be reduced by a finite number of braid moves to the normal form

$$[(12),(12),(13),(13),\ldots,(1\,d-1)(1\,d-1);1,1,\ldots,1,(1d)] \tag{14}$$

where each $(1i)$ appears twice for $1 \leq i \leq d-1$. Without loss of generality we may assume that $(t_1,\ldots,t_n;\lambda_1,\mu_1,\ldots,\lambda_g,\mu_g)$ has minimal $(|\underline{\lambda}|,|\underline{\mu}|)$ and maximal $\ell(\underline{\Sigma})$ as in 4.4 (*)

Step 1. *We claim* $\lambda_1 = \cdots = \lambda_g = 1$. Let us first suppose that $\lambda_1 \neq 1$. Let (ab) be a transposition such that $|(ab)\lambda_1| < |\lambda_1|$. Let $s = [\lambda_1,\mu_1]\cdots[\lambda_g,\mu_g]$. First suppose that $s \neq 1$. Since $s \in A_d$ we have $|s| \geq 2$, thus $n + |s| \geq 2d-2$. By Lemma 4.3 $\langle t_1,\ldots,t_n\rangle$ is a transitive group, so by Mochizuki's Lemma 3.3 we can replace $(t_1,\ldots,t_n)$ by a braid-equivalent sequence $(t'_1,\ldots,t'_n)$ such that $t'_1 = (ab)$. Applying τ''_{11} (cf. Eq.(10)) one transforms λ_1 into $(ab)\lambda_1$. This contradicts the minimality of $(|\underline{\lambda}|,|\underline{\mu}|)$. Suppose that $s = 1$. Then according to Lemma 4.3 the subgroup $\langle t_1,\ldots,t_n\rangle$ has two orbits: $\underline{\Sigma} = \{\Sigma_1,\Sigma_2\}$ where $\#\Sigma_1 = d-1$, $\#\Sigma_2 = 1$. Moreover replacing if necessary $(t_1,\ldots,t_n)$ by a braid-equivalent sequence we may assume that $t_i = t_{i+1}$ for every $i \equiv 1 (mod\, 2)$. Varying over all Hurwitz systems with $\lambda_1, \mu_1, \ldots, \lambda_g, \mu_g$ fixed and braid-equivalent to the given one we want to figure out which transpositions may appear at the first place. Let $\Sigma_1 = \{a_1, \ldots, a_{d-1}\}$, $\Sigma_d = \{a_d\}$. If $\{\alpha, \beta\} \subseteq \Sigma_1$ then by Mochizuki's Lemma 3.3 one may replace the sequence $(t_1, \ldots, t_n)$ by a braid-equivalent one $((a_\alpha, a_\beta), \ldots)$. Let $G_1 = \langle t_1, \ldots, t_n \rangle$, $G_2 = \langle \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g \rangle$. By hypothesis $G = \langle G_1, G_2 \rangle = S_d$ so $\exists h \in G_2$ such that ${\Sigma_1}^h \nsubseteq \Sigma_1$. Let ${\Sigma_1}^h = \{1, \ldots, d\} - \{a_c\}$, so ${a_d}^h = a_c$ with $c < d$. By Main Lemma 2.1 the Hurwitz system $({t_1}^h, \ldots, {t_n}^h; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$ is braid-equivalent to the given one. Applying again Mochizuki's Lemma 3.3 we see that every $(a_\alpha b_\beta)$ with $(\alpha, \beta) \neq (c, d)$ may be placed first in some braid-equivalent Hurwitz system. We may vary $h \in G_2$ with the property ${\Sigma_1}^h \nsubseteq \Sigma_1$, or equivalently with the property ${a_d}^h \in \Sigma_1$. If the orbit ${a_d}^{G_2}$ has ≥ 3 elements, then an arbitrary transposition τ may be placed first in some braid-equivalent Hurwitz system $(t'_1, \ldots, t'_n; \lambda_1, \mu_1, \ldots, \lambda_g, \mu_g)$. Doing this for $\tau = (ab)$ and applying the braid move τ''_{11} we transform λ_1 into $(ab)\lambda_1$ thus obtaining a contradiction with the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$. It remains to consider the case $\# {a_d}^{G_2} = 2$. In this case $G_2 \subseteq S(\{a_c, a_d\}) \times S(\Sigma_1 - \{a_c\})$. If $\lambda_1 \in S(\Sigma_1 - \{a_c\})$ or $\lambda_1 = (a_c a_d)\lambda'_1$ with $\lambda'_1 \in S(\Sigma_1 - \{a_c\})$, $\lambda'_1 \neq 1$ we might decrease $|\lambda_1|$ transforming λ_1 into $(ab)\lambda_1$ with an appropriate (ab) as we saw above and this contradicts the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$. It remains to consider the case $\lambda_1 = (a_c a_d)$. Let $e \in \{1, \ldots, d-1\}$, $e \neq c$. Replacing the n-tuple $(t_1, \ldots, t_n)$ by a braid-equivalent one we may assume that it equals $((a_c a_e), (a_c a_e), \ldots)$. Since $G_2 \subseteq S(\{a_c, a_d\}) \times S(\Sigma_1 - \{a_c\})$ we have $[\lambda_1, \mu_1] = 1$. So $\rho''_{11} : t_1 \mapsto \lambda_1 t_1 \lambda_1^{-1}$. Applying this braid move we obtain $((a_e a_d), (a_c a_e), \ldots; \lambda_1, \mu'_1, \ldots)$. We may replace $(a_e a_d), (a_c a_e)$ by the braid-equivalent pair $(a_c a_d), (a_e a_d)$ and then apply τ''_{11} transforming λ_1 into $(a_c a_d) \cdot (a_c a_d) = 1$. This contradicts again the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$. This proves $\lambda_1 = 1$. If $\lambda_1 = \cdots = \lambda_{k-1} = 1$, but $\lambda_k \neq 1$ we have $u_{k-1} = [\lambda_1, \mu_1] \cdots [\lambda_{k-1}, \mu_{k-1}] = 1$. The braid move τ''_{1k} transforms λ_k into $t_1^{-1}\lambda_k$ according to Theorem 1.8 so the same argument as above may be applied proving that $\lambda_k \neq 1$ contradicts the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$. Therefore $\lambda_1 = \cdots = \lambda_g = 1$.

Step 2. *We claim the group $G_1 = \langle t_1, \ldots, t_n \rangle$ has orbits $\Sigma_1 = \{a_1, \ldots, a_{d-1}\}$, $\Sigma_2 = \{a_d\}$ and there exists $c \in [1, d-1]$ such that every μ_ℓ equals either 1 or $(a_c a_d)$.* For every Hurwitz system with $\lambda_1 = \cdots = \lambda_g = 1$ we have by Theorem 1.8 (b) that ρ'_{1k} transforms μ_k into $t_1^{-1}\mu_k$. Assume that $\mu_1 = \cdots = \mu_{k-1} = 1$ and $\mu_k \neq 1$. Upon a substitution $\tau \leftrightarrow \rho$ we may use arguments similar to those of Step 1 trying to decrease $|\mu_k|$ and thus obtaining a contradiction with the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$. We are not allowed however to use braid moves of the types τ'_{ik} and τ''_{ik} since these change λ_k while we want to preserve at every braid move the equality $\lambda_1 = \cdots = \lambda_g = 1$. We thus have the following: $G_1 = \langle t_1, \ldots, t_n \rangle$ has orbits $\Sigma_1 = \{a_1, \ldots, a_{d-1}\}$, $\Sigma_2 = \{a_d\}$, $\mu_k = (a_c a_d)$, $G_2 = \langle \mu_k, \ldots, \mu_g \rangle \subseteq S(\{a_c, a_d\}) \times S(\Sigma_1 - \{a_c\})$. Here if $\ell > k$ we have either $\mu_\ell = \mu'_\ell \in S(\Sigma_1 - \{a_c\})$ or $\mu_\ell = (a_c a_d)\mu''_\ell$ with $\mu''_\ell \in S(\Sigma_1 - \{a_c\})$. If $\mu'_\ell \neq 1$ or $\mu''_\ell \neq 1$ respectively we might apply Mochizuki's Lemma 3.3 and the braid move $\rho'_{1\ell}$ in order to decrease $|\mu_\ell|$. This is impossible by the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$ so for every $\ell > k$ either $\mu_\ell = 1$ or $\mu_\ell = (a_c a_d)$

Step 3. *We claim $\mu_\ell = 1$ for $\forall \ell < g$, $\mu_g = (a_c a_d)$.* Since renumbering and braid moves are commutative operations we may assume without loss of generality that $\Sigma_1 = \{1, \ldots, d-1\}$ and $c = 1$ so every μ_ℓ equals either 1 or $(1d)$. Let k be the minimal index such that $\mu_k = (1d)$.

If $k = g$ there is nothing to prove, so suppose that $k < g$. Replacing eventually $(t_1, \dots, t_n)$ by a braid-equivalent sequence we may assume that $(t_1, \dots, t_n; \lambda_1, \mu_1, \dots, \lambda_g, \mu_g)$ has the following form:

$$((12),(12),(13),(13),\dots,(1\,d-1)(1\,d-1);1,1,\dots,1,\underset{k}{(1d)},\dots). \tag{15}$$

First suppose that $\mu_g = 1$. We claim that replacing μ_g by $(1d)$ one obtains a braid-equivalent Hurwitz system. Indeed τ'_{1k} (cf. Corollary 1.9) transforms Eq.(15) into $((2d),(12),\dots;$ $\dots,(2d),(1d),\dots)$. Apply the following sequence of braid-equivalences: replace $(2d),(12)$ by $(12),(1d)$; move this pair to the end by braid moves of type σ'_j; perform ρ''_{ng}; move the pair $(12),(1d)$ backward to the front by braid moves of type σ''_j; replace it by $(2d),(12)$ and finally apply τ''_{1k}. One obtains

$$((12),(12),(13),(13),\dots,(1\,d-1)(1\,d-1);1,1,\dots,1,\underset{k}{(1d)},\dots,1,(1d)) \tag{16}$$

as claimed. We may thus suppose that Eq.(15) equals Eq.(16). Applying τ'_{1g} to Eq.(16) and replacing $(2d),(12)$ by $(1d),(2d)$ we obtain

$$((1d),(2d),(13),(13),\dots;\dots,1,(1d),\dots,(2d),(1d))$$

Applying ρ''_{1k} one transforms $\mu'_k = (1d)$ into 1 without changing $t_1 = (1d)$. Replacing $(1d),(2d)$ by $(2d),(12)$ and applying τ''_{1g} we obtain a Hurwitz system with $\lambda'_1 = \dots = \lambda'_g = 1$, $\mu'_1 = \dots = \mu'_k = 1$ and $\mu'_\ell = 1$ or $(1d)$ for $\ell > k$. This shows that the assumption $\mu_k \neq 1$ for $k < g$ contradicts the minimality of $(|\underline{\lambda}|, |\underline{\mu}|)$.

We have so far worked with Hurwitz systems. When working with equivalence classes we may moreover renumber arbitrarily $\{1, \dots, d\}$. We thus conclude that the equivalence class (modulo inner automorphisms) of a Hurwitz system of the type of Step 3 is braid-equivalent to the equivalence class Eq.(15) with $k = g$, so it has the normal form Eq.(14) as claimed ☐

5 The case $n = 2d - 6,\ g = 1$

Unless otherwise specified in this section we shall work with Hurwitz systems $(t_1, \dots, t_n; \lambda, \mu)$ where $n \geq 2$ and $t_1, \dots, t_n$ are transpositions in S_d.

Proposition 5.1. *Let* $(t_1, \dots, t_n; \lambda, \mu)$, *with* $\lambda = 1$, *be a Hurwitz system with monodromy group* S_d. *Suppose that* $d - 1 \leq n < 2d - 2$. *Let* $e = \frac{n}{2} + 1$. *Then the equivalence class* $[t_1, \dots, t_n; \lambda, \mu]$ *is braid-equivalent to*

$$[(12),(12),(13),(13),\dots,(1e),(1e);1,(1\,e+1\dots d)] \tag{17}$$

Proof. We may assume without loss of generality that $(|\lambda|, |\mu|)$ is minimal and $\ell(\underline{\Sigma})$ is maximal in lexicographic order as in 4.4 (*). By Lemma 4.3 the subgroup $\langle t_1, \dots, t_n \rangle \subseteq S_d$ has a single nontrivial domain of transitivity Σ_1 and furthermore since $t_1 \cdots t_n = 1$ we may assume that $t_i = t_{i+1}$ for $i \equiv 1 (mod\, 2)$. Let $\Sigma_1 = \{a_1, \dots, a_d\}$. Decomposing μ into a product of independent cycles $\mu = \mu_1 \cdots \mu_k$ we have each μ_i contains at most one element of Σ_1. Indeed, otherwise using Mochizuki's Lemma 3.3 and applying the braid move ρ''_{ng} (cf. Eq.(10)) we might decrease $|\mu|$. Furthermore the transitivity of $\langle t_1, \dots, t_n, \mu \rangle$ implies that every μ_i contains exactly one element of Σ_1 and each element of $\overline{\Sigma_1}$ is contained in one of μ_i. The inequality $n \geq d-1$, equivalent to $e > d-e$, implies that at least one element of Σ_1 does not appear in the cycles μ_i, $1 \leq i \leq k$. We claim that if $k \geq 2$ then there is a braid-equivalent Hurwitz system $(t'_1, \dots, t'_n; 1, \mu')$ such that μ' is a cycle of order $d-e+1$ containing a single element of Σ_1. Braid moves and renumbering of $\{1, \dots, d\}$ are commutative operations, so for proving the claim we may assume without loss of generality that $(t_1, \dots, t_n; 1, \mu)$ equals

$$((12), (12), \dots, (1e), (1e); 1, (2\ e+1 \dots e+i_1)(3\ e+i_1+1 \dots) \dots). \tag{18}$$

Performing τ'_{11} we obtain

$$((1\, e+1), (12), \dots, (1e), (1e); (1\, e+1), (2\, e+1 \dots)(3 \dots) \dots).$$

Moving the pair $(1\, e+1), (12)$ to the end, replacing it by $(12), (2\, e+1)$ and performing ρ''_{n1} we obtain

$$((13), (13), \dots, (1e), (1e), (12), (12); (1\, e+1), (2\, e+2 \dots)(3\, e+i_1+1 \dots) \dots). \tag{19}$$

Let us consider the system

$$((12), (12), (13), (13), \dots, (1e), (1e); 1, (2\ e+2 \dots e+i_1)(3\ e+1\ e+i_1+1 \dots) \dots). \tag{20}$$

Moving the pair $(13), (13)$ to the front and then performing τ'_{11} we obtain

$$((1\, e+1), (13), (12), (12), (14), (14), \dots; (1\, e+1), (2\, e+2 \dots)(3\, e+1 \dots) \dots).$$

Moving $(1\, e+1), (13)$ to the end, replacing it by $(13), (3\, e+1)$ and performing ρ''_{n1} we obtain

$$((12), (12), (14), \dots, (1e), (13), (13); (1\ e+1), (2\ e+2 \dots)(3\ e+i_1+1 \dots) \dots). \tag{21}$$

Since Eq.(20) is braid-equivalent to Eq.(21) and Eq.(21) is braid-equivalent to Eq.(19) we conclude that Eq.(18) is braid-equivalent to Eq.(20). The sequence Eq.(20) is obtained from Eq.(18) by moving one element from the cycle μ_1 to the cycle μ_2. Repeating this transformation several times we obtain a cycle μ' of order $d-e+1$. We thus proved that replacing the given Hurwitz system by a braid-equivalent one we may assume that $\langle t_1, \dots, t_n \rangle$ has a single nontrivial orbit $\Sigma_1 = \{a_1, \dots, a_e\}$ and $\mu = (a_i a_{e+1} \dots a_d)$ where $1 \leq i \leq e$ and $\{a_1, a_2, \dots, a_d\} = \{1, \dots, d\}$. Renumbering and normalizing the sequence $(t_1, \dots, t_n)$ according to the classical result of Clebsch and Hurwitz we obtain that $[t_1, \dots, t_n; 1, \mu]$ is braid-equivalent to the equivalence class Eq.(17). □

Remark 1. The proof of Proposition 5.1 may be easily adapted in order to show that every equivalence class $[t_1, \dots, t_n; \lambda, 1]$ with full monodromy group S_d and $d-1 \le n < 2d-2$ is braid-equivalent to

$$[(12), (12), \dots, (1e), (1e); (1\,e+1\dots d), 1]. \tag{22}$$

Here one should modify Condition 4.4(*) assuming without loss of generality $(|\lambda|, |\mu|) = (|\lambda|, 0)$ is minimal in the reverse lexicographical order and $\ell(\underline{\Sigma})$ is maximal in lexicographical order. One then repeats the proof of the proposition using ρ'_{n1} instead of τ'_{11} and τ''_{11} instead of ρ''_{n1}.

Lemma 5.2. *Let $(t_1, \dots, t_n; \lambda, \mu)$ be a Hurwitz system with monodromy group S_d. Let $\underline{\Sigma} = \{\Sigma_1, \dots, \Sigma_m\}$ be the domains of transitivity of $\langle t_1, \dots, t_n\rangle$ ordered in such a way that $\#\Sigma_1 \ge \#\Sigma_2 \ge \cdots \ge \#\Sigma_m$. Suppose that $(|\lambda|, |\mu|)$ is minimal and $\ell(\underline{\Sigma})$ is maximal as in 4.4(*). Suppose that $\lambda \ne 1$. Then the following three conditions cannot hold simultaneously.*

i. Σ_1 is non-minimal (cf. 4.2).

ii. There exists $a \in \Sigma_1$ such that $a^\lambda \notin \Sigma_1$ (equivalently there exists $c \in \Sigma_1$ such that $c^{\lambda^{-1}} \notin \Sigma_1$).

iii. There exists $b \in \Sigma_1$ such that $b^\lambda = b$.

Proof. Decomposing λ into a product of independent cycles $\lambda = \lambda_1 \cdots \lambda_k$ we see that each of the two conditions of (ii) is equivalent to the existence of a λ_i which contains elements of both Σ_1 and $\overline{\Sigma_1} = \{1, \dots, d\} - \Sigma_1$. Let $\mu = \mu_1 \cdots \mu_\ell$ be the decomposition of μ into a product of independent cycles. The minimality of $(|\lambda|, |\mu|)$ implies that neither λ_i nor μ_j may contain two different elements of some Σ_r. In fact if this were the case then we might apply Mochizuki's Lemma 3.3 and one of the braid moves τ''_{11} or ρ''_{n1} in order to decrease $(|\lambda|, |\mu|)$. In particular for each $x \in \Sigma_1$ we have an alternative: either $x^\lambda = x$ or $x^\lambda \notin \Sigma_1$.

Suppose, by way of contradiction, that (i), (ii) and (iii) hold simultaneously. Let $s = [\lambda, \mu]$. By Lemma 4.3 we must have $n + |s| < 2d-2$ since otherwise $\langle t_1, \dots, t_n\rangle$ is transitive, so applying Mochizuki's Lemma 3.3 and the braid move τ''_{11} we might decrease $|\lambda|$ which contradicts the minimality of $(|\lambda|, |\mu|)$ assumed in the lemma. Lemma 4.3 yields moreover that $n_1 + |s|_{\Sigma_1}| = 2(\#\Sigma_1 - 1)$, so Condition (i) means $s|_{\Sigma_1}$ has $q \ge 2$ domains of transitivity $\Delta_1, \dots, \Delta_q$. We claim one can choose a and b satisfying (ii) and (iii) respectively so that they belong to different domains. In fact if $b \in \Delta_k$ and all $x \in \Sigma_1$ with $x^\lambda \notin \Sigma_1$ belong to Δ_k then we may replace b by an arbitrary element of some other Δ_ℓ according to the alternative of the preceding paragraph. Let $a \in \Delta_i,\ b \in \Delta_j,\ i \ne j$. Let T_1 be the subsequence of $T = (t_1, \dots, t_n)$ consisting of the transpositions that move elements of Σ_1. Let $a' = a^s,\ b' = b^s$. Using Corollary 3.6 we can replace T_1 by a sequence $(\cdots, (a'b'), (a'b'))$ and then move the pair $(a'b'), (a'b')$ to the end of T. We then perform the braid move ρ'_{n1} which transforms $t_n = (a'b')$ into $(a'b')^{s^{-1}\lambda} = (ab)^\lambda = (a^\lambda b)$. The group generated by the new sequence $\langle T'\rangle$ has orbit $\Sigma_1 \cup \{a^\lambda\}$. By Mochizuki's Lemma 3.3 one may replace T' by $((a\,a^\lambda), \dots)$. Performing τ''_{11} one transforms λ into $(a\,a^\lambda)\lambda$ and clearly $|(a\,a^\lambda)\lambda| < |\lambda|$. This contradicts the minimality of $(|\lambda|, |\mu|)$. □

Lemma 5.3. *Let* $(t_1,\dots,t_n;\lambda,\mu)$ *be a Hurwitz system with monodromy group* S_d*. Let* $d-1\le n<2d-2$*. Suppose that* $(|\lambda|,|\mu|)$ *is minimal and* $\ell(\underline{\Sigma})$ *is maximal as in 4.4(*). Suppose that* $s=[\lambda,\mu]=1$*. Then* $\lambda=1$ *and the equivalence class* $[t_1,\dots,t_n;\lambda,\mu]$ *is braid-equivalent to Eq.*(17)*.*

Proof. Let $\underline{\Sigma}=\{\Sigma_1,\dots,\Sigma_m\}$ be the orbits of $\langle t_1,\dots,t_n\rangle$. Let $e=\frac{n}{2}+1$. According to Lemma 4.3 we have $\#\,\Sigma_i=1$ for each $i\ge 2$ and $\#\,\Sigma_1=e$. Suppose, by way of contradiction, that $\lambda\ne 1$. According to Lemma 5.2 its conditions (i) – (iii) cannot hold simultaneously. The orbit Σ_1 is non-minimal since $s|_{\Sigma_1}=1$. As we saw in the course of the proof of Lemma 5.2 the minimality of $(|\lambda|,|\mu|)$ implies that if one decomposes λ into a product of independent cycles $\lambda=\prod\lambda_i$ then every λ_i contains at most one element of Σ_1. The inequality $n\ge d-1$ implies that $\#\,\Sigma_1=e>d-e$. Thus there exists an element $b\in\Sigma_1$ such that $b^\lambda=b$. Therefore Condition (ii) of Lemma 5.2 must fail to hold, i.e. we have $a^\lambda=a$ for every $a\in\Sigma_1$. Let $F\subseteq\{1,\dots,d\}$ be the set of fixed points of λ. We have $F\supset\Sigma_1$ so the group $\langle t_1,\dots,t_n\rangle$ leaves F invariant. Since $[\lambda,\mu]=1$ we have $F^\mu=F$. Therefore the assumption $\lambda\ne 1$ implies that $\langle t_1,\dots,t_n,\lambda,\mu\rangle$ is not a transitive group which is a contradiction. The last part of the lemma refers to Proposition 5.1. ☐

Remark 2. This lemma may be stated differently: if $[\lambda,\mu]=1$ and $\lambda\ne 1$ then one can decrease $|\lambda|$ by a sequence of braid moves (at the expense of possible increasing of $|\mu|$). Modifying appropriately Lemma 5.2 one can prove similarly that given a Hurwitz system $(t_1,\dots,t_n;\lambda,\mu)$ with full monodromy group S_d and such that $d-1\le n<2d-2$, then $[\lambda,\mu]=1$ and $\mu\ne 1$ imply that one can decrease $|\mu|$ by a sequence of braid moves (at the expense of possible increasing of $|\lambda|$).

Theorem 5.4. *Let* $d\ge 4$, $n=2d-6$*. Let* Y *be an elliptic curve. Then the Hurwitz space* $\mathcal{H}^0_{d,n}(Y)$ *parametrizing simple coverings with monodromy groups equal to* S_d *is irreducible.*

Proof. By Theorem 1.10 it suffices to show that given a Hurwitz system $(t_1,\dots,t_n;\lambda,\mu)$ with monodromy group S_d its equivalence class is braid-equivalent to

$$[(12),(12),(13),(13),\dots,(1\,d-2),(1\,d-2);1,(1\,d-1\,d)]. \tag{23}$$

Without loss of generality we may assume that $(t_1,\dots,t_n;\lambda,\mu)$ has minimal $(|\lambda|,|\mu|)$ and maximal $\ell(\underline{\Sigma})$ as in 4.4(*). Here $\underline{\Sigma}=\{\Sigma_1,\dots,\Sigma_m\}$ are the domains of transitivity of $G_1=\langle t_1,\dots,t_n\rangle$ ordered in such a way that $\#\,\Sigma_1\ge\#\,\Sigma_2\ge\cdots\ge\#\,\Sigma_m$. We aim to prove that $\lambda=1$ in order to apply Proposition 5.1. Let $s=[\lambda,\mu]$. Our first goal is to prove $s=1$. Suppose, by way of contradiction, that $s\ne 1$, so in particular $\lambda\ne 1$ and $\mu\ne 1$. Since $[\lambda,\mu]$ is an even permutation we have $|s|\equiv 0(mod 2)$. If $s\ge 4$ then $n+|s|\ge 2d-2$, so by Lemma 4.3 the group $G_1=\langle t_1,\dots,t_n\rangle$ is transitive. Using Mochizuki's Lemma 3.3 and the braid move τ''_{11} we can transform λ into λ' with $|\lambda'|<|\lambda|$. This contradicts the minimality of $(|\lambda|,|\mu|)$. Thus if $s\ne 1$ then $|s|=2$. By Lemma 4.3 the group G_1 has two domains of transitivity $\underline{\Sigma}=\{\Sigma_1,\Sigma_2\}$ and Σ_2 is minimal. If $\lambda=\lambda_1\cdots\lambda_k,\ \mu=\mu_1\cdots\mu_\ell$ are the factorizations into independent cycles, then as we saw in the proof of Lemma 5.2 the minimality of $(|\lambda|,|\mu|)$

implies that each λ_i or μ_j is a transposition of the type (ac) where $a \in \Sigma_1,\ c \in \Sigma_2$. It is clear that $\max\{k, \ell\} \leq \#\,\Sigma_2$. Let $s_1 = (s|_{\Sigma_1})^{\sim},\ s_2 = (s|_{\Sigma_2})^{\sim}$ where $^{\sim}$ means trivial extension from $S(\Sigma_i)$ to S_d. We have $s = s_1 s_2$ and the following cases may occur: a) $|s_1| = 2,\ |s_2| = 0;$ b) $|s_1| = 1,\ |s_2| = 1;$ c) $|s_1| = 0,\ |s_2| = 2$.

In Case (a) by minimality $\#\,\Sigma_2 = 1$. So $\lambda_1 = (a_1 c),\ \mu = (a_2 c)$ where $\{c\} = \Sigma_2$ and $a_1 \neq a_2$ since $[\lambda, \mu] \neq 1$. Then $[\lambda, \mu] = (a_2 a_1 c)$. This contradicts $[\lambda, \mu] = t_1 \cdots t_n \in S(\Sigma_1) \times S(\Sigma_2)$, so Case (a) is impossible.

In Case (b) we have $\#\,\Sigma_2 = 2$ by minimality and $\#\,\Sigma_1 \geq 2$. If $d \geq 5$ then $\#\,\Sigma_1 \geq 3$. So Σ_1 is non-minimal and furthermore there exists an element $b \in \Sigma_1$ such that $b^{\lambda} = b$. If $\lambda = \lambda_1$ or $\lambda = \lambda_1\lambda_2$ with $\lambda_1 = (ac)$ then $a \in \Sigma_1,\ a^{\lambda} \notin \Sigma_1$. This contradicts Lemma 5.2. It remains to consider the case $d = 4$. Let $\Sigma_1 = \{a_1, a_2\},\ \Sigma_2 = \{c_1, c_2\}$.

Suppose that λ is a transposition. We may assume that $\lambda = (a_1 c_1)$. We exclude case by case the possibilities for μ. The cases $\mu = (a_1 c_1)$, or $(a_2 c_2)$, or $\mu = (a_1 c_1)(a_2 c_2)$ are impossible since $[\lambda, \mu] \neq 1$. The cases $\mu = (a_1 c_2)$ or $\mu = (a_2 c_1)$ are impossible either since $[\lambda, \mu] = s_1 s_2$ is not a cycle of order 3. It remains to exclude the possibility $\mu = (a_1 c_2)(a_2 c_1)$. Indeed here

$$[\lambda, \mu] = (a_1 c_1)(a_1 c_2)(a_2 c_1)(a_1 c_1)(a_1 c_2)(a_2 c_1) = (a_1 c_1)(a_2 c_2),$$

while $[\lambda, \mu]$ must equal $(a_1 a_2)(c_1 c_2)$.

Suppose that $\lambda = \lambda_1\lambda_2$. We may assume that $\lambda = (a_1 c_1)(a_2 c_2)$. Then $[\lambda, \mu]$ might be $\neq 1$ only if $\mu = (a_1 c_2)$ or $\mu = (a_2 c_1)$. The calculation above shows that in these cases $[\lambda, \mu] \neq (a_1 a_2)(c_1 c_2)$ So Case (b) is impossible.

In Case (c) we have $\#\,\Sigma_2 = 3$ by minimality and $s|_{\Sigma_1} = 1$. Here $d = \#\,\Sigma_1 + \#\,\Sigma_2 \geq 6$. If $d \geq 7$ we obtain a contradiction with Lemma 5.2 by the same argument we used in Case (b). The cases $d = 6$ and $\lambda = \lambda_1$ or $\lambda = \lambda_1\lambda_2$ are excluded in the same way. It remains to consider the case $\lambda = (a_1 c_1)(a_2 c_2)(a_3 c_3)$ where $\Sigma_1 = \{a_1, a_2, a_3\},\ \Sigma_2 = \{c_1, c_2, c_3\}$. Let T_i be the subsequence of $T = (t_1, \ldots, t_6)$ which moves elements of $\Sigma_i,\ i = 1, 2$. Replacing T by a braid-equivalent sequence we may assume that $T = T_1 T_2$ where $T_1 = (a_1 a_2)(a_1 a_2)(a_1 a_3)(a_1 a_3)$. By Main Lemma 2.1 the Hurwitz system $(T_1 T_2; \lambda, \mu)$ is braid-equivalent to $({T_1}^{\mu} T_2; \lambda, \mu)$. If $\#({\Sigma_1}^{\mu} \cap \Sigma_2) < 3$ then the group $\langle {T_1}^{\mu}, T_2\rangle$ has a domain of transitivity $\Sigma_1' = {\Sigma_1}^{\mu} \cup \Sigma_2$ and $\#\Sigma_1' > \#\Sigma_1$. This contradicts the maximality of $\ell(\underline{\Sigma})$. Hence ${\Sigma_1}^{\mu} = \Sigma_2,\ {\Sigma_2}^{\mu} = \Sigma_1$. This implies that the group $(t_1, \ldots, t_n; \lambda, \mu)$ is imprimitive which is excluded by hypothesis. We thus excluded all possible cases with $s \neq 1$. Therefore $s = [\lambda, \mu] = 1$.

Suppose that $d \geq 5$. Then from Lemma 5.3 and Proposition 5.1 it follows that $\lambda = 1$ and $[t_1, \ldots, t_n; \lambda, \mu]$ is braid-equivalent to Eq.(23) which proves the theorem when $d \geq 5$. So, the only case that remains to be considered is $d = 4, n = 2, [\lambda, \mu] = 1$. Here we have $\underline{\Sigma} = \{\Sigma_1, \Sigma_2, \Sigma_3\}$ with $\#\Sigma_1 = 2,\ \#\,\Sigma_2 = \#\,\Sigma_3 = 1$. Let $\overline{\Sigma_1} = \Sigma_2 \cup \Sigma_3$. Suppose, by way of contradiction, that $\lambda \neq 1$. Unless $\lambda = (ac)(bd)$ where $\Sigma_1 = \{a, b\},\ \overline{\Sigma_1} = \{c, d\}$ we may apply the argument of Lemma 5.3 and obtain a contradiction with the minimality of $(|\lambda|, |\mu|)$. We obtain a Hurwitz system of the type $((ab), (ab); (ac)(bd), \mu)$ where μ commutes with $(ac)(bd)$. The centralizer of $\lambda = (ac)(bd)$ is $\{1, (ac)(bd), (ad)(bc), (ab)(cd), (ac), (bd), (abcd), (adcb)\}$. The braid move ρ''_{21} transforms μ into $\mu'' = (ab)\mu$ (cf. Eq.(10)). Thus the minimality of $(|\lambda|, |\mu|)$ excludes the cases $\mu = (ab)(cd), (abcd)$ or $(adcb)$. The hypothesis of primitivity of $\langle t_1, t_2, \lambda, \mu\rangle$

excludes the cases $\mu = 1, (ac)(bd)$ or $(ad)(bc)$. It thus remains to consider the cases $\mu = (ac)$ or $\mu = (bd)$. These are equivalent up to reordering, so it suffices to consider $\mu = (ac)$. We perform the following sequence of braid moves starting from $((ab),(ab);(ac)(bd),(ac))$. Applying τ'_{11} we obtain $((bc),(ab);(acdb),(ac))$. Replacing $(bc),(ab)$ by $(ac),(bc)$ and then applying τ''_{11} (cf. Theorem 1.8 (f)) we obtain $(ac) \mapsto (ac)^{(abc)} = (ab)$ and $(acdb) \mapsto (ac)(acdb) = (adb)$. So we obtain $((ab),(bc);(bad),(ac))$. Applying τ''_{11} again we transform $(\dots;(bad),(ac))$ into $(\dots;(bd),(ac))$. This contradicts the minimality of $(|\lambda|,|\mu|)$ of the initial Hurwitz system $((ab),(ab);(ac)(bd),(ac))$. All possible cases with $[\lambda,\mu] = 1,\ \lambda \neq 1$ were excluded, so $\lambda = 1$. We conclude that $(t_1,t_2;\lambda,\mu) = ((ab),(ab);1,\mu)$. The minimality of $(|\lambda|,|\mu|) = (0,|\mu|)$ implies that no independent cycle of the factorization of μ may contain both a and b. Since $\langle (ab),\mu\rangle$ is transitive we conclude that the factorization is either $\mu = \mu_1\mu_2 = (ac)(bd)$ with $\{b,d\} = \overline{\Sigma}_1$ or μ is a cycle of order 3 containing the two elements of $\overline{\Sigma}_1$. The former case is impossible since $\langle (ab),\mu\rangle$ is a primitive group. We conclude that the equivalence class $[(ab),(ab);1,\mu]$ equals $[(12),(12);1,(134)]$. The theorem is proved. □

We can sharpen Theorem 3.14 using the same proof and Theorem 3.10, Theorem 4.5 and Theorem 5.4.

Theorem 5.5. *Let Y be a smooth, projective curve of genus ≥ 1. Let $d \geq 2,\ n > 0,\ n \equiv 0 (mod\ 2)$. If d is prime then $\mathcal{H}_{d,n}(Y) = \mathcal{H}^0_{d,n}(Y)$. If d is not prime let d' be its maximal divisor $\neq d$. Suppose that $n \geq \max(2, 2d'-4)$ or if $g = 1$ suppose that $n \geq \max(2, 2d'-6)$. Then the connected components of $\mathcal{H}_{d,n}(Y)$ which correspond to simple coverings with monodromy groups $\neq S_d$ are in one-to-one correspondence with the equivalence classes of unramified coverings $[\tilde{Y} \to Y]$ of degrees $d_2|d$ where $d_2 \neq 1, d$.*

References

[Bi] Joan S. Birman, *On braid groups*, Comm. Pure Appl. Math. **22** (1968), 41–72.

[Cl] Clebsch A., *Zür Theorie der Riemann'schen Fläche*, Math. Ann. **6** (1872), 216–230.

[FB] Edward Fadell and James Van Buskirk, *The braid groups of E^2 and S^2*, Duke Math. J. **29** (1962), 243–257.

[FN] Edward Fadell and Lee Neuwirth, *Configuration spaces*, Math. Scand. **10** (1962), 111–118.

[Fu] William Fulton, *Hurwitz schemes and irreducibility of moduli of algebraic curves*, Ann. of Math. (2) **90** (1969), 542–575.

[GHS] Graber T., Harris J. and Starr J., *A note on Hurwitz schemes of covers of a positive genus curve*, preprint, arXiv:math.AG/0205056.

[Hu] Hurwitz A., *Ueber Riemann'schen Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. **39** (1891), 1–61.

[Kl] Paul Kluitmann, *Hurwitz action and finite quotients of braid groups*, Braids (Santa Cruz, CA, 1986) (Providence, RI), Contemp. Math., vol. 78, Amer. Math. Soc., Providence, RI, 1988, pp. 299–325.

[Lo] Eduard Looijenga, *On the tautological ring of* $\mathcal{M}_g$, Invent. Math. **121** (1995), no. 2, 411–419.

[Mo] Shinichi Mochizuki, *The geometry of the compactification of the Hurwitz scheme*, Publ. Res. Inst. Math. Sci. **31** (1995), no. 3, 355–441.

[Na] S. M. Natanzon, *Topology of two-dimensional coverings, and meromorphic functions on real and complex algebraic curves. I* (in Russian), Trudy Sem. Vektor. Tenzor. Anal. (1988), no. 23, 79–103. (English translation: Selecta Math. Soviet. **12** (1993), no. 3, 251–291.)

[Sc] G. P. Scott, *Braid groups and the group of homeomorphisms of a surface*, Proc. Cambridge Philos. Soc. **68** (1970), 605–617.

[Ve] Vetro F., *Irreducibility of Hurwitz spaces of coverings of elliptic curve of prime degree with one point of total ramification*, Matematiche (Catania) (to appear), Preprint N. 192, May 2003, Dipartimento di Matematica, Università di Palermo.

[Vo] Helmut Völklein, *Groups as Galois groups*, Cambridge Studies in Advanced Mathematics, vol. 53, Cambridge University Press, Cambridge, 1996.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PALERMO
VIA ARCHIRAFI N.34, 90123 PALERMO, ITALY and
INSTITUTE OF MATHEMATICS, BULGARIAN ACADEMY OF SCIENCES

E-mail address: **kanev@math.unipa.it**